arXiv:1510.08135v2 [math.AT] 4 Apr 2016

ALGEBRAIC COBORDISM AND FLAG VARIETIES

N.YAGITA

Abstract. Let X be an algebraic variety over k such that $\bar{X} = X \otimes_k \bar{k}$ is cellular. We study torsion elements in the Chow ring $CH^*(X)$ which corresponds to v_iy in the algebraic cobordism $\Omega^*(\bar{X})$ where $0 \neq y \in CH^*(\bar{X})/p$ and v_i is the generator of BP^* with $|v_i| = -2(p^i - 1)$. In particular, we compute $CH^*(X)$ for the twisted complete flag variety $X = \mathbb{G}_k/B_k$ over k.

1. Introduction

Let X be a smooth algebraic variety over a field k of $ch(k) = 0$ such that $\bar{X} = X \otimes_k \bar{k}$ is cellular. For a fixed prime p, let $CH^*(X) = CH^*(X)_{(p)}$ be the Chow ring of algebraic cycles modulo rational equivalence, localized at p, and $\Omega^*(X)$ the BP^*-version of the algebraic cobordism $MGL^{2*,*}(X) \otimes_{MU^*} BP^*$ defined by Voevodsky and Levine-Morel ([Vo1], [Le-Mo1,2], [Ya2,6]), with the coefficient ring $\Omega^* = BP^* \cong \mathbb{Z}_{(p)}[v_1, v_2, ...]$, where $|v_n| = -2(p^n - 1)$. The relation between these theories are given ([Vo1],[Le-Mo1,2])

$$CH^*(X) \cong \Omega^*(X) \otimes_{\Omega^*} \mathbb{Z}_{(p)} \cong \Omega^*(X)/(\Omega^{<0}).$$

In this paper, we define an invariant $deg_v(y)$ to study the restriction image $res_\Omega : \Omega^*(X) \to \Omega^*(\bar{X})$. Let F^{2i} be the Ω^*-subalgebra of $\Omega^*(\bar{X})$ generated by elements in $CH^j(\bar{X})$ for $j \geq i$. Let $gr^*\Omega(\bar{X}) = \oplus_i F^i/F^{i+1}$ be the associated graded algebra. Let us write

$$Res_\Omega = gr^*(Image(res_\Omega)) \subset gr^*(\Omega^*(\bar{X})).$$

Let $y \neq 0 \in CH^*(\bar{X})/p$ (namely, y is also identified with an Ω^*-module generator of $\Omega^*(\bar{X})$). If $v_iy \in Res_\Omega$, then we can easily see that $v_jy \in Res_\Omega$ for all $j \leq i$, by using the Landweber-Novikov operations. So we define $deg_v(y) = n$ if $v_{n-1}y \in Res_\Omega$ but $v_ny \notin Res_\Omega$. (Let $deg_v(y) = 0$ if $y \in Res_\Omega$ and $deg_v(y) = -1$ if $py \notin Res_\Omega$.)

If $deg_v(y) = n > 0$, then there are nonzero element $c_i(y)$ in $CH^*(X)$ for $0 \leq i \leq n-1$, which corresponds to v_iy in $\Omega^*(\bar{X})$ generating Res_Ω as an Ω^*-module (since $y \notin Res_\Omega$).

2000 *Mathematics Subject Classification.* 55N20, 55R12, 55R40.

Key words and phrases. Algebraic cobordism, Rost motive, flag manifold.

Lemma 1.1. *If $deg_v(y) = n > 0$, then there are $c_i = c_i(y) \in CH^*(X)$ for $0 \leq i \leq n-1$ with $|c_i| = |y| - 2(p^i - 1)$ such that the following $M_n(y)$ is contained (as a quotient) in an irreducible motive $CH^*(M)$ in $CH^*(X)$,*

$$M_n(y) = \mathbb{Z}_{(p)}\{c_0\} \oplus \mathbb{Z}/p\{c_1, ..., c_{n-1}\} \twoheadleftarrow CH^*(M) \twoheadleftarrow CH^*(X).$$

(Here the notation $A\{a, b, ...\}$ means the free A-module generated by $a, b, ...$)

When $deg_v(y) \leq 0$, let $M_0(y) = \mathbb{Z}\{y\}$ and $M_{-1}(y) = 0$.

Corollary 1.2. *Let $CH^*(\bar{X}) \cong \oplus_i \mathbb{Z}_{(p)}\{y_i\}$. Then $CH^*(X)$ has the quotient module such that*

$$Q(X) = \oplus_i M_{d_i}(y_i) \twoheadleftarrow CH^*(X) \quad \textit{where } d_i = deg_v(y_i).$$

In this paper, we study $CH^*(X)$ for a non-algebraically closed field k by using this $deg_v(y)$. In general, to compute $deg_v(y)$ is a quite difficult problem. However there are cases $deg_v(y)$ are known or computable.

Given a pure symbol a in the mod p Milnor K-theory $K^M_{n+1}(k)/p$, by Rost, we can construct the norm variety V_a of $dim(V_a) = p^n - 1$ (such that $\bar{V}_a$ is cellular). Rost and Voevodsky showed that there is $y \in CH^{b_n}(\bar{V}_a)$ for $b_n = (p^n - 1)/(p - 1)$ such that y^{p-1} is the fundamental class of $\bar{V}_a$. The results by Rost and Voevodsky show that $deg_v(y^i) = n$ for all $1 \leq i \leq p - 1$, and moreover we know ([Ro1], [Vo1,4], [Me-Su], [Vi-Ya], [Ya6])

Theorem 1.3. *For a nonzero pure symbol $a \in K^M_{n+1}(k)/p$, there is an irreducible (Rost) motive R_a (write by R_n simply) in the motive $M(V_a)$ of the norm variety V_a such that $CH^*(\bar{R}_a) \cong \mathbb{Z}_{(p)}[y]/(y^p)$, and*

$$CH^*(R_a) \cong \mathbb{Z}_{(p)} \oplus \oplus_{i=1}^{p-1} M_n(y^i) \cong \mathbb{Z}_{(p)} \oplus M_n(y)[y]/(y^{p-1}).$$

Let G be a simply connected compact Lie group (and T its maximal torus) and G_k be the corresponding split reductive group over k and T_k its split torus. Let $\mathbb{G}_k$ be a nontrivial G_k-torsor, and $\mathbb{G}_k/T_k$ the (twisted form of) complete flag variety. We are interesting in the Chow ring $CH^*(\mathbb{G}_k/T_k)$. (For a Borel subgroup $B \supset T$, note G_k/B_k is cellular and $CH^*(\mathbb{G}_k/B_k) \cong CH^*(\mathbb{G}_k/T_k)$.)

We only consider here the cases

$$grH^*(G; \mathbb{Z}/p) \cong \mathbb{Z}/p[y]/(y^p) \otimes \Lambda(x_1, ..., x_\ell)$$

where the degree are $|y| = even$, and $|x_i| = odd$ (for example, exceptional Lie groups except for $(G, p) = (E_8, 3)$). We call these group G of type (I). There is a fibering $G \to G/T \to BT$ and the induced spectral sequence (see for details, [Tod]. [To-Wa], [Mi-Ni], [Na])

$$E_2^{*,*'} \cong H^*(G; \mathbb{Z}/p) \otimes S(t) \Longrightarrow H^*(G/T; \mathbb{Z}/p)$$

where $S(t) = H^*(BT) \cong \mathbb{Z}_{(p)}[t_1, ..., t_\ell]$ with $|t_i| = 2$. Then we can write

$$grH^*(G/T; \mathbb{Z}/p) \cong \mathbb{Z}/p[y]/(y^p) \otimes S(t)/(\bar{b}_1, ..., \bar{b}_\ell)$$

where $d_{|x_i|+1}(x_i) = \bar{b}_i$ for the differential in the spectral sequence.

Petrov, Semenov and Zainoulline [Pe-Se-Za] prove that when G is a group of type (I), the motive $M(X)$ (which is localized at p) is decomposed as a sum of (degree changing) Rost motive R_2, namely,

$$CH^*(X)/p \cong CH^*(R_2)/p \otimes S(t)/(\bar{b}_1, ..., \bar{b}_\ell)$$

On the other hand, the following additive isomorphism is immediate

$$S(t)/(p, \bar{b}_i\bar{b}_j, \bar{b}_k) \cong \mathbb{Z}/p\{1, \bar{b}_1, ..., \bar{b}_{2p-2}\} \otimes S(t)/(\bar{b}_1, ..., \bar{b}_\ell)$$

where i, j, k range $1 \leq i, j \leq 2p-2 < k \leq \ell$.

Theorem 1.4. *Let G be a simply connected simple Lie group of type (I). Let $X = \mathbb{G}_k/T_k$ for a nontrivial torsor $\mathbb{G}_k$. Then $deg_v(y^r) = 2$ and $c_s(y^r) = b_{2r-s} \in S(t)$ with $b_j = \bar{b}_j$ mod(p) where $s = 0$ or 1, and $1 \leq 2r, j \leq 2(p-1)$. Hence we have the ring isomorphism*

$$CH^*(X)/p \cong S(t)/(p, \bar{b}_i\bar{b}_j, \bar{b}_k).$$

where i, j, k range $1 \leq i, j \leq 2(p-1) < k \leq \ell$.

The following corollaries are proved for $G = G_2$ and $p = 2$ in [Ya5,7].

Corollary 1.5. *Let G be a group in the above theorem. Then we have the isomorphism $CH^*(\mathbb{G}_k)/p \cong \mathbb{Z}/p$.*

Let $gr_\gamma(\bar{X})$ be the graded ring associated by the γ-filtration of $K^0(\bar{X})_{(p)}$. (See §15 below for the γ-filtration.)

Corollary 1.6. *Let G be a group in the above theorem. Then we have the isomorphism $gr_\gamma(\bar{X}) \cong gr_\gamma(X) \cong CH^*(X)$.*

After the first version of this paper was written, Karpenko [Kar] showed that when $\mathbb{G}_k$ is generically twisted, then $S(t) \to CH^*(X)$ is always surjective. Hence above two corollaries are just corollaries of Karpenko's general result. However some advantage of ours are that we can get ring structure of $gr_\gamma(X)$ form Theorem 1.4.

This paper is organized as follows. In §2 we recall the motivic BP-theory using the Atiyah-Hirzebruch spectral sequence. In §3, we define $deg_v(y)$ and study it for the norm variety in §4. In §5, we study the relation between $deg_v(y)$ and the Dickson invariant. We recall the Chow ring for the split (or topological) flag manifold $X = G/T$ in §6, and begin to study it for non split cases in §7. In §8 we recall the results by Petrov, Semenov and Zainoulline. In §9, we study the type (I) groups, and show the main result. In §10, we write down the

case $G = G_2, F_4$ explicitly. In §11 we consider the cases where G is a products of type (I) groups, and §12 is a very short note for a type (II) group. In §13, we study the case when X are anisotropic quadrics, and in §14, we study the case $G = SO(m)$. In the last section, we study the γ filtration in the (algebraic and topological) K-theories.

2. ABP theories

Recall that $MU^*(-)$ is the complex cobordism theory defined in the usual (topological) spaces and

$$MU^* = MU^*(pt.) \cong \mathbb{Z}[x_1, x_2, ...] \quad |x_i| = -2i.$$

Here each x_i is represented by a sum of hyper surfaces of $dim(x_i) = 2i$ in some product of complex projective spaces. Let $MGL^{*,*'}(-)$ be the motivic cobordism theory defined by Voevodsky [Vo1]. Let us write by AMU the spectrum $MGL_{(p)}$ in the stable $\mathbb{A}^1$-category representing the motivic cobordism theory (localized at p), i.e.,

$$MGL^{*,*'}(-)_{(p)} = AMU^{*,*'}(-).$$

Here note that $AMU^{2*,*}(pt.) \cong MU^{2*}_{(p)}$ (but is not isomorphic to $AMU^{*,*'}(pt)$, in general). In particular $AMU^{*,*'}(X)$ is an $MU^*_{(p)}$-algebra.

Given a regular sequence $S_n = (s_1, ..., s_n)$ with $s_i \in MU^*_{(p)}$, we can inductively construct the AMU-module spectrum by the cofibering of spectra ([Ya2,4])

$$(2.1) \quad \mathbb{T}^{-1/2|s_i|} \wedge AMU(S_{i-1}) \xrightarrow{\times s_i} AMU(S_{i-1}) \to AMU(S_i)$$

where $\mathbb{T} = \mathbb{A}/(\mathbb{A} - \{0\})$ is the Tate object. For the realization map $t_{\mathbb{C}}$ induced from $k \subset \mathbb{C}$, it is also immediate that $t_{\mathbb{C}}(AMU(S_n)) \cong MU(S_n)$ with $MU(S_n)^* = MU^*/(S_n)$.

Recall that the Brown-Peterson cohomology theory $BP^*(-)$ with the coefficient $BP^* \cong \mathbb{Z}_{(p)}[v_1, v_2...]$ by identifying $v_i = x_{p^i-1}$. (So $|v_i| = -2(p^i - 1)$.) We can construct spectra

$$ABP = AMU(x_i | i \neq p^j - 1)$$

such that $t_{\mathbb{C}}(ABP) \cong BP$. For $S = (v_{i_1}, ..., v_{i_n})$, let us write

$$ABP(S) = AMU(S \cup \{x_i | i \neq p^j - 1\})$$

so that $t_{\mathbb{C}}(ABP(S)) = BP(S)$ with $BP(S)^* = BP^*/(S)$.

In particular, let $AH\mathbb{Z} = ABP(v_1, v_2, ...)$ so that $AH\mathbb{Z}^{2*,*}(pt.) \cong \mathbb{Z}_{(p)}$. In the $\mathbb{A}^1$-stable homotopy category, Hopkins-Morel showed that

$$AH\mathbb{Z} \cong H_{\mathbb{Z}} \quad , i.e., \quad AH\mathbb{Z}^{*,*'}(X) \cong H^{*,*'}(X, \mathbb{Z}_{(p)})$$

the (usual) motivic cohomology. Using this result, we can construct the motivic Atiyah-Hirzebruch spectral sequence.

Theorem 2.1. *([Ya2,6]) Let $Ah = ABP(S)$ for $S = (v_{i_1}, v_{i_2}, ...)$. Then there is AHss (the Atiyah-Hirzebruch spectral sequence)*

$$E(Ah)_2^{(*,*',2*'')} = H^{*,*'}(X; h^{2*''}) \Longrightarrow Ah^{*+2*'',*'+*''}(X)$$

with the differential $d_{2r+1} : E_{2r+1}^{(*,*',2*'')} \to E_{2r+1}^{(*+2r+1,*'-r,2*''-2r)}$.

Note that the cohomology $H^{m,n}(X, h^{2n'})$ here is the usual motivic cohomology with coefficients in the abelian group $h^{2n'}$. Here we recall some important properties of the motivic cohomology. When X is smooth, we know

$$CH^*(X) \cong H^{2*,*}(X; \mathbb{Z}_{(p)}), \quad H^{*,*'}(X; \mathbb{Z}_{(p)}) \cong 0 \ for \ 2*' < *.$$

Hence if X is smooth, then $E_r^{m,n,2n'} \cong 0$ for $m > 2n$. The convergence in AHss means that there is the filtration

$$Ah^{*,*'}(X) = F_0^{*,*'} \supset F_1^{*,*'} \supset F_2^{*,*'} \supset ...$$

such that $F_i^{*,*'}/F_{i+1}^{*,*'} \cong E_\infty^{*+2i,*'+i,-2i}$. Let $S \subset R = (v_{j_1}, ...)$. Then the induced map $ABP(S) \to ABP(R)$ of spectra induces the $BP_{(p)}^*$-module map of AHss : $E(ABP(S))_r^{*,*',*''} \to E(ABP(R))_r^{*,*',*''}$.

From the above notes and dimensional reason, we see

$$ABP(S)^{2*,*}(pt) \cong BP(S)^* = BP^*/(S). \tag{2.3}$$

We also have for a smooth X,

$$ABP(S)^{2*,*}(X) \otimes_{BP^*} \mathbb{Z}_{(p)} \cong H^{2*,*}(X) \cong CH^*(X). \tag{2.4}$$

More generally, we have

Lemma 2.2. *Let X be smooth and $S \subset R \subset (p, v_1, ...)$. Then*

$$ABP(R)^{2*,*}(X) \cong ABP(S)^{2*,*}(X)/(R).$$

Remark. Note that in general degree $(*, *')$,

$$ABP(R)^{*,*'}(X) \ncong ABP(S)^{*,*'}(X)/(R).$$

In this paper, we mainly consider $ABP^{2*,*}(X)$. So we write it simply

$$\Omega^{2*}(X) = ABP^{2*,*}(X),$$

which is isomorphic to

$$MGL^{2*,*}(X) \otimes_{MU^*} BP^* \cong \Omega^*{}_{LM}(X) \otimes_{MU^*} BP^*$$

for the algebraic cobordism $\Omega^*{}_{LM}(X)$ define by Levine and Morel.

Let y_i generate (as an Ω^*-module) $\Omega^*(X)$ (hence $y_i \neq 0 \in CH^*(X)/p$). We consider the graded ring given by degree of $|y_i|$ (two times of degree in $CH^*(X)$), i.e.

$$gr\Omega^*(\bar{X}) = \oplus F^i/F^{i+1}, \quad for\ F^i = \sum_{|y_j|\geq i} \Omega^*(y_j).$$

(In this paper, we use the notations that $\Omega^*(y)$ means the Ω^*-module generated by y, and $\Omega^*\{y\}$ means the Ω^*-free module generated by y.) Hence we note that

$$gr\Omega^{2*}(X) \cong \oplus_{*''} E_\infty^{2*,*,*''}$$

for the infinity term of the AHss converging to $ABP^{*,*'}(X)$.

Recall that $I_n = (p = v_0, v_1, ..., v_{n-1})$ and $I_\infty = (v_0, v_1, ...)$ are the prime invariant ideals (under Landweber-Novikov operations) of BP^*.

Lemma 2.3. *Let x be an Ω^*-module generator of $\Omega^*(X)$. If $v_n x = 0$ in $gr\Omega^*(X)$, then $v_i x = 0$ in $gr\Omega^*(X)$ for all $0 \leq i \leq n$ (i.e., $I_{n+1}x = 0$).*

Proof. Recall the Quillen (Landweber-Novikov) operation r_α in the $\Omega^*(-)$ theory ([Ha], [Ra], [Ya2,6]) such that

$$r_{p^i\Delta_{n-i}}(v_n) = v_i \mod(I_i^2).$$

Let $\alpha = p^i\Delta_{n-i}$. Then by the Cartan formula, we have

$$r_\alpha(v_n x) = r_\alpha(v_n)\cdot x + \sum_{\alpha'+\alpha''=\alpha,\ |\alpha''|>0} r_{\alpha'}(v_n) r_{\alpha''}(x).$$

Here $r_{\alpha''}(x) \in F_{|r_{\alpha''}(x)|} \subset F_{|x|+2}$. Hence in $gr\Omega^*(X)$ we get $r_\alpha(v_n x) = (v_i + a_i)x$ with $a_i \in I_i^2$. By induction on i, we see if $v_n x = 0 \in gr\Omega^*(X)$, then so is $v_i x$ for $0 \leq i \leq n$. □

Definition. Let x be an Ω^*-module generator of $\Omega^*(X)$ (or $\Omega^*(M)$ for a motive M in the motive $M(X)$ of X). Define the v-torsion type of x by $tor_v(x) = m$ if $v_m x \neq 0$ but $v_{m-1}x = 0$ in $gr\Omega^*(X)$. Moreover we define

$$tor_v(X) = max_x tor_v(x)$$

where x ranges Ω^*-module generators of $\Omega^*(X)$.

Example. If X is cellular, then $tor_v(X) = 0$. Let R_a be the Rost motive defined from a nonzero symbol $0 \neq a \in K^M_{n+1}(k)/p$. Then $tor_v(R_a) = n$ but $tor_v(\bar{R}_a) = 0$ where $\bar{R}_a = R_a \otimes \bar{k}$ for the algebraically closed field $\bar{k}$. (See §4 for details,)

Proposition 2.4. *Let X be smooth. If $dim(X) \leq 2(p^n - 1)$, then $tor_v(X) \leq n$.*

Proof. Let $A\tilde{k}(n)^{*,*'}(X)$ be the motivic connected integral Morava K-theory with the coefficient

$$A\tilde{k}(n)^{2*,*}(pt.) \cong \tilde{k}(n)^* = \mathbb{Z}_{(p)}[v_n].$$

We consider the following two AHss's

$$E(BP)_2^{*,*',*''} \cong H^{*,*'}(X; BP^{*''}) \Longrightarrow ABP^{*,*'}(X),$$

$$E(\tilde{k}(n))_2^{*,*',*''} \cong H^{*,*'}(X; \tilde{k}(n)) \Longrightarrow A\tilde{k}(n)^{*,*'}(X),$$

and induced map $\rho : E(BP)_r^{*,*',*''} \to E(\tilde{k}(n))_r^{*,*',*''}$.

Since $|v_n| = -2(p^n - 1)$, we see

$$E(\tilde{k}(n))_r^{*,*',*''} = 0 \quad unless\ *'' = 0\ mod(2(p^n - 1)).$$

Recall degree of the differential $deg(d_{2r+1}) = (2r+1, -r, -2r)$. Hence $d_{2r+1} = 0$ unless $r = 0\ mod(p^n - 1)$. Hence if $dim(X) \leq (p^n - 1)$, then all differential are zero, and the spectral sequence collapses, i.e., $E(\tilde{k}(n))_2 \cong E(\tilde{k}(n))_\infty$. In particular $v_n x \neq 0$ in $E(\tilde{k}(n))_\infty$ for all $0 \neq x \in CH^*(X)$.

Suppose $tor_v(x') \geq n+1$ for some $0 \neq x' \in CH^*(X)$. Then $v_n x' = 0$ $mod(I_\infty^2)$ in $gr\Omega(X) \cong E(BP)_\infty$. This contradicts to the compatibility of the map ρ of AHss's. □

Remark. Vishik showed more strong results. He proves (Theorem 4.3 in [Vi]) that as an Ω^*-module, $\Omega^*(X)$ has relations only of positive degree. If $tor_v(y) \geq n+1$, then there is a relation $v_n y = 0$ in $gr\Omega^*(X)$ (and hence relation $v_n y + v_{n+1} y' + ... = 0$ in $\Omega^*(X)$). So $|y| > -|v_n| = 2(p^n - 1)$.

3. Res

Let us write $\bar{X} = X \otimes \bar{k}$ for the algebraically closure $\bar{k}$ of k, and $i_{\bar{k}} : \bar{X} \to X$ the induced map. For an (oriented) cohomology theory $h^*(X)$, let us write the image of the restriction map

$$Res_h = grIm(i_{\bar{k}}^*) \subset grh^*(\bar{X})$$

for $res_h = i_{\bar{k}}^* : h^*(X) \to h^*(\bar{X})$.

Lemma 3.1. *Let $y \in \Omega^*(\bar{X})$ be an Ω^*-module generator.*
If $v_n y \in Res_\Omega$, then $v_j y \in Res_\Omega$ for all $0 \leq j \leq n$, namely,

$$I_{n+1}\{y\} \subset Res_\Omega \subset gr\Omega^*(\bar{X}).$$

Proof. The proof is quite similar to that of Lemma 2.3. Recall the Quillen (Landweber-Novikov) operation r_α in the $\Omega^*(-)$ theory. By the arguments similar to the proof of Lemma 2.3, we see that $r_\alpha(v_n y) =$

$(v_i + a)y \in Res_\Omega$ for $a \in I_i^2$. By induction on i, we can see $v_i y \in Res_\Omega$ for all $0 \leq n$. □

Definition. For a smooth X and an Ω^*-module generator $y \in \Omega^*(\bar{X})$ with $y \notin Res_\Omega$, we say that y has v-degree n

$$deg_v(y) = n \quad if\ v_{n-1}y \in \mathrm{Res}_\Omega\ but\ v_n y \notin Res_\Omega.$$

When $y \in Res_\Omega$, let $deg_v(y) = 0$ and when $py \notin Res_{CH}$ let $deg_v(y) = -1$. Moreover define

$$deg_v(X) = max_{0 \neq y \in CH^*(\bar{X})/p} deg_v(y).$$

Lemma 3.2. *If M is an irreducible Ω^*-module, then M is contained in one irreducible motive in $\Omega^*(X)$.*

Proof. Each map from $\Omega^*(X)$ to $\Omega^*(Y)$ of motives is represented by a composition map $f_* g^*$ of the induced map g^* (of smooth varieties) and the Gysin map f_*. The both maps are Ω^* maps. Hence a decomposition of an Ω^*-motive gives that of an Ω^*-module. □

Hereafter in this section, we assume the following condition for y and X such that we can define $deg_v(y)$.

(1) $\Omega^*(\bar{X})$ is an Ω^*-free module.
(2) $y \in \Omega^*(\bar{X})$ with $0 \neq y \in CH^*(\bar{X})/p$.

Lemma 3.3. *If $deg_v y = n > 0$, then there are $c_i \in CH^*(X)$ for $0 \leq i \leq n-1$ with $|c_i| = |y| - 2(p^i - 1)$ such that the following $M_n(y)$ is contained (as a quotient) in an irreducible motive $CH^*(M)$ in $CH^*(X)$,*

$$M_n(y) = \mathbb{Z}_{(p)}\{c_0\} \oplus \mathbb{Z}/p\{c_1, ..., c_{n-1}\} \twoheadleftarrow CH^*(M) \twoheadleftarrow CH^*(X).$$

Proof. We recall that

$$CH^*(X) \cong \Omega^*(X) \otimes_{BP^*} \mathbb{Z}_{(p)} \cong \Omega^*(X)/(BP^{<0}\Omega^*(X))$$

where $BP^{<0} = (v_1, ..., v_n, ...)$ is the ideal of BP^* generated by negative degree elements. The assumption $deg_v(y) = n$ implies $I_n\{y\} \subset \Omega^*\{y\} \cap Res_\Omega$. Moreover since $v_j y \notin Res_\Omega$ for all $j \geq n$, we see

$$I_n\{y\} = \Omega^*\{y\} \cap Res_\Omega \mod ((BP^{<0})^2\Omega^*\{y\}).$$

Here we note

$$I_n/((BP^{<0})^2 \cap I_n) \cong (p, ..., v_{n-1})/((BP^{<0}) \cdot (p, ..., v_{n-1}))$$
$$\cong \mathbb{Z}_{(p)}\{p\} \oplus \mathbb{Z}/p\{v_1, ..., v_{n-1}\} \cong M_n.$$

Since $p \cdot v_i = v_i \cdot p \in BP^{<0}I_n$ for $i > 0$, we have $pv_i = 0$ in M_n. Hence we see

$$M_n = (Res_\Omega \cap \Omega^*\{y\})/(BP^{<0}Res_\Omega) \cap \Omega^*\{y\}) \twoheadleftarrow (res_\Omega(CH^*(X))_{(p)}).$$

(Note that there is a projection $\Omega^*(\bar{X}) \twoheadrightarrow \Omega^*\{y\}$ since $\Omega^*(\bar{X})$ is Ω^*-free.) Thus there is a surjection $CH^*(X) \twoheadrightarrow M_n(y)$.

Let us consider the decompositions of motives

$$\Omega^*(X) \cong \oplus_i \Omega^*(M_i), \quad \Omega^*(\bar{X}) \cong \oplus_{i,j} \Omega^*(N_{i,j})$$

with $\Omega^*(\bar{M_i}) \cong \oplus_j \Omega^*(N_{i,j})$. Take (s,t) such that $\Omega^*\{y\} \subset \Omega^*(N_{s,t})$. Then

$$M_n(y) \twoheadleftarrow \Omega^*(X)/(\oplus_{j \neq s} \Omega^*(M_j)) \cong \Omega^*(M_s).$$

This shows that $M_n(y)$ is contained in one irreducible motive. □

Corollary 3.4. *Let X split over a field k' over k of index $p \cdot a$ for some a coprime to p. Then for all $0 \neq y \in CH^*(\bar{X})/p$, we see*
(1) $py \in Res$.
(2) for $i \neq 0$, $pc_i(y) = 0$, i.e.,
additively $M_n(y) \subset CH^*(X)$ for $deg_v(y) = n$.
(3) If $dim(X) \leq (p^n - 1)$, then $0 \leq deg_v(X) \leq n$.

Proof. For $i > 0$, we see $res_{CH}(c_i) = 0$. By using transfer, we have (2)

$$pc_i = tr \cdot res(c_i) = tr(0) = 0.$$

We will show $py \in Res_{CH}$ for this case. Using the fact that $res \otimes \mathbb{Q}$ is isomorphic, there is s such that $p^s y = res(x)$ for some $x \in CH^*(X)$. Then

$$p^s res \cdot tr(y) = res \cdot tr \cdot res(x) = res(apx) = ap^{s+1}(y).$$

Since $CH^*(\bar{X})$ is torsion free, we have $res \cdot tr(a^{-1}y) = py$.

If $deg_v(y) \geq n+1$, then there exists non zero element c_n and hence $|c_n| = |y| - 2(p^n - 1)$ must be positive. □

Corollary 3.5. *Assume that $res_\Omega : \Omega^*(X) \to \Omega^*(\bar{X})$ is injective. If $deg_v(y) = n > 0$, then $tor_v(c_i) = i$ for each $0 \leq i \leq n-1$.*

Proof. Let $0 \leq i < j \leq n-1$. Of course in $\Omega^*(\bar{X})$, $v_i v_j y = v_j v_i y$. The injectivity of res_Ω implies that $v_i c_j = v_j c_i$ in $\Omega^*(X)$. Since $|c_j| < |c_i|$, we get $v_i c_j = 0$ in $gr\Omega^*(X)$. Hence $tor_v(c_j) \geq j$.

On the other hand, suppose that $v_j c_j = 0$ in $gr\Omega^*(X)$. Then there is a relation in $\Omega^*(X)$ such that

$$v_j c_j - \sum_s w_s d_s = 0 \quad for\ w_s \in \Omega^*,\ |v_j| > |w_s|.$$

This implies that $v_j^2 y = \sum w_s res_\Omega(d_s)$ in $\Omega^*(\bar{X})$ which is Ω^*-free. Note $w_s \neq v_j$ since $|w_s| < |v_j|$. This means that $w_s = v_j^2$, and $y = res_\Omega(d_s)$ which contradicts to $deg_v(y) > 0$. □

Remark. For each invariant ideal J (under Landweber-Novikov operations) in BP^*, we can do the similar arguments. For example if $J\{y\} = Res_\Omega \cap \Omega^*\{y\}$ for $J = (p, v_1^2)$, then $deg_v(y) = 1$ but we have

$$M(y)' = \mathbb{Z}\{c_0\} \oplus \mathbb{Z}/p\{c_1'\} \twoheadleftarrow CH^*(X)$$

where $c_1' = v_1^2 y$ and $|c_1'| = |y| - 4(p^2-1)$. For the case $J = (p^2, pv_1, v_1^2)$. we have

$$M(y)'' = \mathbb{Z}\{c_{0,0}\} \oplus \mathbb{Z}/p\{c_{0,1}, c_{1,1}\} \twoheadleftarrow CH^*(X)$$

with $c_{0,0} = p^2 y, c_{0,1} = pv_1 y, c_{1,1} = v_1^2 y$. This case my appear for the product of $X \times X$ for $deg_v(X) = 2$. (See §11 below.)

Remark. When the assumption (1) before Lemma 3.3 does not hold, this lemma does not hold. Let $tor_v(y) = m > 0$ and $deg_v(y) = n$, and $m < n$. By elements $v_m y, ..., v_{n-1} y$, we can define element $c_m, ..., c_{n-1} \in CH^*(X)$. However we do not have the quotient

$$M_{(m,n)}(y) = \mathbb{Z}/p\{c_m, ..., c_{n-1}\} \twoheadleftarrow CH^*(X).$$

In fact, suppose that there is a relation $v_i y' + v_j y = 0 \in \Omega^*(\bar{X})$ for $i < j$, and $m \le j \le n-1$. If $y' \in Res_{CH}$ but $y \notin Res_{CH}$, then

$$c_j = c_j(y) = v_j y = -v_i y' = 0 \in \Omega^*(X) \otimes_{\Omega^*} \mathbb{Z}_{(p)} \cong CH^*(X).$$

Recall $M_n(y)$ is defined for $n > 0$ in the previous lemma. Moreover, let $M_0(y) = \mathbb{Z}_{(p)}\{y\}$ and $M_{-1}(y) = 0$ (when $py \notin Res_\Omega$).

Corollary 3.6. *Let $CH^*(\bar{X}) \cong \oplus_i \mathbb{Z}_{(p)}\{y_i\}$. Then $CH^*(X)$ has the quotient module such that*

$$Q(X) = \oplus_i M_{d_i}(y_i) \twoheadleftarrow CH^*(X) \quad where\ d_i = deg_v(y_i).$$

Corollary 3.7. *For $n \ge 0$, let $f : X \to Y$ be a map of smooth varieties such that*

$$f^* : CH^*(\bar{Y})/p \to CH^*(\bar{X})/p \quad is\ injective\ for\ \ * \le n,$$

and that if $deg_v(y) \ge 1$ then $deg(f^*(y)) \ge 1$ for all $0 \ne y \in CH^*(\bar{Y})/p$ with $* \le n$. Then $deg_v(y) \le deg_v(f^*(y))$ and

$$f^* : Q^*(Y) \to Q^*(X) \quad is\ injective\ for\ \ * \le n.$$

Proof. If $v_j y_i \in Res_{\Omega^*}$, then $f^*(v_j y_i) = v_j f^*(y_i) \in Res_{\Omega^*}$. Hence $deg_v(y_i) \le deg_v(f^*(y_i))$. For $d \le d'$, it is immediate $M_d \subset M_{d'}$ by the definition. From Corollary 3.6, we have this corollary. □

4. Norm variety

Let X be a smooth (quasi projective) variety. Recall that $H^{*,*'}(X;\mathbb{Z}/p)$ be the $mod(p)$ motivic cohomology defined by Voevodsky and Suslin. Rost and Voevodsky solved the Beilinson-Lichtenbaum (Bloch-Kato) conjecture ([Vo2.4], [Su-Jo], [Ro2])

$$H^{*,*'}(X;\mathbb{Z}/p) \cong H^*_{et}(X;\mu_p^{\otimes *'}) \quad for * \leq *'.$$

In this paper, we always assume that k has a primitive p-th $root\ of\ unity$ (so $\mu_p \cong \mathbb{Z}/p$). Then we have the isomorphism $H^m_{et}(X;\mu_p^{\otimes n}) \cong H^m_{et}(X;\mathbb{Z}/p)$. Let τ be a generator of $H^{0,1}(Spec(k);\mathbb{Z}/p) \cong \mathbb{Z}/p$, so that $colim_i \tau^i H^{*,*'}(X;\mathbb{Z}/p) \cong H^*_{et}(X;\mathbb{Z}/p)$.

Let χ_X be the $\hat{C}ech$ complex, which is defined as $(\chi_X)^n = X^{n+1}$ (see details [Vo1,2,4]). Let $\tilde{\chi}_X$ be defied as a cofiber

$$\tilde{\chi}_X \to \chi_X \to Spec(k)$$

in the stable $\mathbb{A}^1$-homotopy category. Voevodsky defined the motivic cohomology $H^{*,*'}(\chi;\mathbb{Z}/p)$ for all object χ in the stable $\mathbb{A}^1$-homotopy category. Morover, he showed that there exists the Milnor operation in the motivic cohomology ([Vo1,3])

$$Q_i : H^{*,*'}(\chi;\mathbb{Z}/p) \to H^{*+2p^i-1,*'+p^i-1}(\chi;\mathbb{Z}/p)$$

which is compatible with the usual Milnor operation on $H^*(t_{\mathbb{C}}(\chi);\mathbb{Z}/p)$ for the realization map $t_{\mathbb{C}}$. (Here the topological operation is defined $Q_0 = \beta$ Bockstein operation and $Q_{i+1} = Q_i P^{p^i} - P^{p^i} Q_i$.) This operation Q_i can be extended on $H^*(M;\mathbb{Z}/p)$ for a motive M in $M(X)$ (Lemma 7.1 in [Ya]).

For $0 \neq x \in H^{*,*'}(X;\mathbb{Z}/p)$ (or cohomology operation), let us write $* = |x|, w(x) = 2*' - *$ so that $0 \leq |x| \leq 2dim(X)$ and $w(x) \geq 0$ when X is smooth. We also note

$$|\tau| = 0,\ w(\tau) = 2, \quad |Q_i| = 2p^i - 1,\ w(Q_i) = -1.$$

Let $f_X \in CH^{dim(X)}(\bar{X})$ be the fundamental class of $\bar{X}$. Suppose $deg_v(f_X) = n$. It is known that the ideal

$$I(X) = \pi_*(X) \subset BP^*(pt.) = BP^* \quad for\ \pi : X \to pt.$$

is an invariant ideal (e.g. Lemma 5.3 in [Ya]). Since $\pi_*(f_X) = 1$, we get $\pi_*(v_{n-1} f_X) = v_{n-1}$, and this means $I(X) \supset I_n$.

We consider the following assumption for an existence of the motive M in X.

Assumption 4.1. *There is a motive M in $M(X)$ such that*

(1) *there is $y \in CH^*(\bar{M})$ with $deg_v(y) \geq 1$,*

$$(2)\quad H^{*,*'}(M;\mathbb{Z}/p) \cong H^{*,*'}(\chi_X;\mathbb{Z}/p) \quad for\ 0 < *' < * \leq |y|.$$

Lemma 4.2. *Assume that $I(X) \supset I_n$ and (1), (2) in the above assumption. Moreover let $deg_v(y) \geq n$. Then there is $a' \in H^{*,*'}(M;\mathbb{Z}/p)$ such that*

$$c_i(y) = Q_0...\hat{Q}_i...Q_{n-1}(a').$$

Remark. Here $a = c_i(y)$ means only that $res_\Omega(a) = v_i y\ mod(I_\infty^2)$. Of course, $c_i(y)$ is not determined uniquely.

To prove this lemma, we use the following lemmas.

Lemma 4.3. *(Lemma 6.4,6.5 in [Ya]) If $I_n \subset I(X)$, then $ABP^{2*,*}(\tilde{\chi}_X)$ is I_n-torsion, and $H^{*,*'}(\tilde{\chi}_X;\mathbb{Z}/p)$ is $\Lambda(Q_0,...,Q_{n-1})$-free.*

Lemma 4.4. *(Corollary 3.4 in [Ya]) Let $x \in CH^*(X) \cong E_\infty^{2*,*,0}$ and $v_s x = 0$ in $E_\infty^{2*,*,*'}$ for the AHss converging to $ABP^{*,*'}(X)$. Then there is $b \in H^{*,*'}(X;\mathbb{Z}/p)$ with $Q_s(b) = x$ and a relation in $ABP^{2*,*}(X)$*

$$v_s x + v_{s+1}x_{s+1} + ... + v_k x_k + ... = 0 \quad mod(I_\infty^2)$$

with $x_k = Q_k(b)$ in $H^{2*,*}(X;\mathbb{Z}/p)$ for all $k \geq s$.

Proof of Lemma 4.2. Since $deg_v(y) = n$, there is $c_{n-1}(y) \in H^{2*,*}(M;\mathbb{Z}/p)$ which correspond to $v_{n-1}y$. Since $|Q_i c_{n-1}| \leq |y|$ for $i \leq n-2$, we see $Q_i(c_{n-1}) = 0$ from (2) since $w(Q_i c_{n-1}) = -1$. Since $I(X) \supset I_n$, $H^{*,*'}(\tilde{\chi}_X;\mathbb{Z}/p)$ is $\Lambda(Q_0,...,Q_{n-1})$-free from Lemma 4.2. Therefore $c_{n-1}(y)$ is in the Q_i-image. Hence there is $a' \in H^{*,*'}(\tilde{\chi}_X;\mathbb{Z}/p)$ such that

$$Q_0...Q_{n-2}(a') = c_{n-1}(y).$$

Since $c_{n-1}(y) \in Im(Q_0)$, it is a p-torsion in $CH^*(X)$, and this means

$$pc_{n-1}(y) + v_1 c'_{n-2} + ... + v_{n-1}c'_0 = 0 \quad in\ \Omega^*(X)/I_\infty^2$$

for some $c'_i \in \Omega^*(X)$. In particular, we see that

$$res_\Omega(c'_0) = py \in \Omega^*(\bar{X})/(res_\Omega(I_\infty^2)) \cong \Omega^*(\bar{X})/I_\infty^3$$

from $res_\Omega(c_{n-1}(y)) = v_{n-1}y$. (Note $res_\Omega(c'_i) = 0$ for $i \neq 0$.) Thus we can take $c_0(y) = -c'_0$ (note $c_i(y)$ are only decided in $\Omega^*(X)$ modulo $(Ker(res_\Omega), I_\infty)$).

Then from Lemma 4.3, there is $b \in H^{*,*'}(X;\mathbb{Z}/p)$ such that

$$Q_0(b) = c_{n-1}(y), \quad Q_{n-1}(b) = c_0(y).$$

Since $H^{*,*'}(\tilde{\chi}_X;\mathbb{Z}/p)$ is $\Lambda(Q_0,...,Q_{n-1})$-free, this b is determined uniquely, that is $b = Q_1...Q_{n-2}(a')$. Hence we have

$$c_0(y) = Q_{n-1}(b) = Q_{n-1}Q_{n-2}...Q_1(a').$$

We also get $c_i(y) = Q_0...\hat{Q}_i...Q_{n-1}(a')$ by the same argument. □

Let $x = x_0...x_n \in H^{n+1,n+1}(X;\mathbb{Z}/p)$ for $x_i \in H^{1,1}(X;\mathbb{Z}/p)$. Suppose that there are $s \geq 0$ and $x' \in H^{*,*'}(X;\mathbb{Z}/p)$ such that

$$(**) \quad x = \tau^s x' \qquad where\ \tau \in H^{0,1}(Spec(k);\mathbb{Z}/p).$$

Then by dimensional reason such that $w(Q_i) = -1$ and $w(\tau) = 2$,

$$Q_{i_1}...Q_{i_{n+1-2i}}(x') \in CH^*(X)/p,$$

$$(e.g., \quad q_i(s) = Q_0...\hat{Q}_i...Q_{n+1-2s}(x') \in CH^*(X)/p).$$

Since $\bar{X}$ is cellular, we see $H^{odd,*'}(\bar{X};\mathbb{Z}_{(p)}) = 0$. So $res_H(x') = 0$ for $res_H : H^{*,*'}(;\mathbb{Z}/pX) \to H^{*,*'}(\bar{X};\mathbb{Z}/p)$. Therefore $res(q_i(s)) = 0$ also. Since $q_i(s)$ for $i \neq 0$ is in the Q_0-image, it is p-torsion in the integral cohomology. However $q_0(s)$ has a possibility of non-torsion elements.

Lemma 4.5. *Suppose $(**)$ and $I_n \subset I(X)$. Moreover suppose (1),(2) in Assumption 4.1 and $c_0(y) = q_0(s) = Q_1...Q_{n+1-2s}(x')$. Then $deg_v(y) \geq n + 2 - 2s$ and*

$$c_i(y) = q_i(s) = Q_0...\hat{Q}_i...Q_{n+1-2s}(x'). \quad for\ 0 \leq i \leq n+1-2s$$

Proof. Let $b = Q_1...Q_{n-2s}(x')$. Then from Lemma 4.3,4.4, we have the relation

$$pQ_0(b) + v_1Q_1(b) + ... + v_{n+1-2s}Q_{n+1-2s}(b) = 0 \quad in\ \Omega^*(X)/I^2_\infty.$$

Here $Q_i(b) = 0$ for $i \neq 0, n+1-2s$ since $Q_i^2 = 0$. Since $Q_{n+1-2s}(b) = c_0(y)$ by the assumption, we have $res(Q_0(b)) = v_{n+1-2s}y$, namely $Q_0(b) = c_{n+1-2s}(y)$. □

Now we recall the norm variety. Given a pure symbol a in the mod p Milnor K-theory $K^M_{n+1}(k)/p$, by Rost, we can construct the norm variety V_a (such that $\bar{V}_a$ is cellular and) that

$$\pi_*([V_a]) = v_n, \quad a|_{k(V_a)} = 0 \in K^M_{n+1}(k(V_a))$$

where $[V_a] = 1 \in \Omega^0(V_a)$, $\pi : V_a \to pt.$ is the projection and $k(V_a)$ is the function field of V_a over k.

Rost and Voevodsky showed that there is $y \in CH^{b_n}(\bar{V}_a)$ for $b_n = p^{n-1} + ... + p + 1$ such that y^{p-1} is the fundamental class of $\bar{V}_a$, that is

$$\mathbb{Z}_{(p)}\{y^{p-1}\} \cong CH^{p^n-1}(\bar{V}_a) \cong CH_0(\bar{V}_a) \cong \mathbb{Z}_{(p)}.$$

Then $deg(V_a) = p$ and $y^{p-1} \notin Res_{CH}$, but $py^{p-1} \in Res_{CH}$. Moreover the same facts hold for y^i with $1 \leq i \leq p-1$. Hence each y^i satisfies (1),(2) in the preceding (§3) section, in particular $deg_v(y^i) > 0$. More strongly, Rost and Voevodsky show the following theorem.

Theorem 4.6. *([Ro1,2], [Vo1,4]) For a nonzero symbol $a \in K^M_{n+1}(k)/p$, let $M(V_a)$ is the motive of the norm variety V_a. Then there is an irreducible motive R_a (write R_n also) in $M(V_a)$ such that*

$$CH^*(\bar{R}_a) \cong \mathbb{Z}_{(p)}[y]/(y^p)$$

$$CH^*(R_a) \cong \mathbb{Z}_{(p)} \oplus \oplus_{i=1}^{p-1} M_n(y^i) \cong \mathbb{Z}_{(p)} \oplus M_n(y)[y]/(y^{p-1})$$

where $|y| = 2b_n$ and $M_n(y) = \mathbb{Z}_{(p)}\{c_0\} \oplus \mathbb{Z}/p\{c_1, ..., c_{n-1}\}$.

We can also prove

Theorem 4.7. *([Vi-Ya], [Ya6]) The map $res_\Omega : \Omega^*(R_a) \to \Omega^*(\bar{R}_a)$ is injective and*

$$Res_\Omega = \Omega^*\{1\} \oplus I_n[y]^+/(y^p) \subset \Omega^*[y]/(y^p) \cong \Omega^*(\bar{R}_n).$$

Corollary 4.8. *For a Rost motive R_n and $1 \leq i \leq p-1$, we have $deg_v(y^i) = n$ and $tor_v(c_j(y^i)) = j$ for $0 \leq j \leq n-1$.*

By Rost and Voevodsky, it is known that

$$H^{*,*'}(\chi_a; \mathbb{Z}/p) \cong H^{*,*'}(R_a; \mathbb{Z}/p)$$

for all $* \leq 2b_n$, $*' \leq *$, and there is a short exact sequence

$$0 \to H^{*+1,*}(\chi_a; \mathbb{Z}/p) \xrightarrow{\tau} K^M_{*+1}(k)/p \to K^M_{*+1}(k(V_a))/p \to 0$$

where we identify $H^{*+1,*+1}(\chi_a; \mathbb{Z}/p) \cong K^M_{*+1}(k)/p$. Hence there exists $a' = \tau^{-1}(a) \in H^{*+1,*}(R_a; \mathbb{Z}/p)$ ($s = 1$ in $(**)$ and Lemma 4.5).

We can define $y \in CH^*(\bar{R}_a)$ by $q_0(1) = Q_1...Q_{n-1}(a') = c_0(y)$, namely, $res_\Omega(q_0(1)) = py$, and $c_i = q_i(1) = Q_0...\hat{Q}_i...Q_{n-1}(a')$ as in Lemma 4.4. We can prove Theorem 4.6 by using Lemma 4.2, 4.5, while assuming the above facts. Hence the original proof ([Vo1,24]) is better than ours, I write down Lemma 4.2, Lemma 4.5 for possibility of applications for other spaces.

5. Dickson invariant

The contents in this section are not used other sections but arguments are very similar to those given in the preceding section for the element $q_i(s)$.

Throughout this section, we assume that k contains ξ_{p^2} a primitive p^2-root of the unity. Since

$$Q_0(\tau) = \xi_p = 0 \quad in\ k^*/(k^*)^p \cong H^{1,1}(pt.; \mathbb{Z}/p),$$

we see Q_i commutes with τ, i.e., $Q_i(\tau \cdot z) = \tau \cdot Q_i(z)$ for $z \in H^{*,*'}(X; \mathbb{Z}/p)$.

We consider the case that $s = 0$ and $X = B(\mathbb{Z}/p)^n$ (while $\bar{X}$ is of course not cellular) with

$$H^{*,*'}(X; \mathbb{Z}/p) \cong K^M_*(k) \otimes \mathbb{Z}/p[\tau, y_1, ..., y_n] \otimes \Lambda(x_1, ..., x_n)$$

for p odd prime. The graded ring $grH^{*,*'}(X;\mathbb{Z}/2)$ is isomorphic to the right hand side ring (and we consider in this graded ring for $p=2$). Then $x=x_1...x_n$ is invariant under the $SL_n(\mathbb{Z}/p)$-action. So each

$$q_i(0)' = Q_0...\hat{Q}_i...Q_n(x) \quad x=x_1...x_n$$

is $SL_n(\mathbb{Z}/p)$- invariant (the element $q_i(0)$ in the preceding section is defined for $x=x_0x_1...x_n$). The invariant ring on the polynomial ring is well known by Dickson

$$\mathbb{Z}/p[y_1,...,y_n]^{GL_n(\mathbb{Z}/p)} \cong \mathbb{Z}[c_{n,0},....,c_{n,n-1}],$$

$$\mathbb{Z}/p[y_1,...,y_n]^{SL_n(\mathbb{Z}/p)} \cong \mathbb{Z}[e_n,c_{n,1},....,c_{n,n-1}],$$

where $e_n^{p-1}=c_{n,0}$ and each $c_{n,i}$ is defined by

$$\Pi_{y\in\mathbb{Z}/p\{y_1,...,y_n\}}(t+y) = t^{p^n} + \sum t^{p^i}c_{n,i}.$$

These $c_{n,i}$ is also represented by Chern classes. Let $reg:(\mathbb{Z}/p)^n \to U_{p^n}$ be the (complex) regular representation. Then

$$c_{n,i} = c_{p^n-p^i}(reg) = reg^*(c_{p^n-p^i}).$$

It is known by Mimura-Kameko [Mi-Ka]

$$e_nc_{n,i} = Q_0...\hat{Q}_i...Q_n(x_1...x_n).$$

Hence the element $e_nc_{n,i}$ is just $q_i(0)'$. Hence we get

$$q_i(0)' = c_{n,i}e_n = reg^*(c_{p^n-1})^{1/(p-1)}reg^*(c_{p^n-p^i}).$$

Each element $a\in H^{1,1}(X;\mathbb{Z}/p) \cong H^1_{et}(X;\mathbb{Z}/p)$ can be represented by $\mathbb{A}^1$-homotopy map $X\to B\mathbb{Z}/p$. So we can write

$$a=a_1...a_n = i_a^*(x_1...x_n) \quad for\ i_a: X\to (BZ/p)^n.$$

Theorem 5.1. *For an element $a=a_1...a_n\in H^{n,n}(X;\mathbb{Z}/p)$, we have*

$$q_i(0)' = Q_0...\hat{Q}_i...Q_n(a)$$

$$= i_a^*(e_nc_{n,i}) = i_a^*((reg^*(c_{p^n-1}))^{1/(p-1)}reg^*(c_{p^n-p^i})).$$

It is known ([Vo1,2]) that

$$H^{*,*'}(B(\mathbb{Z}/2)^n;\mathbb{Z}/2) \cong K_*^M(k)\otimes\mathbb{Z}/2[\tau,y_1,...,y_n,x_1,...,x_n]/(Re.)$$

$$where\ \ Re. = (x_i^2=\tau y_i \mid 1\le i\le n).$$

Giving grading by multiplying τ, we have

$$(*)\quad grH^{*,*'}(B(\mathbb{Z}/2)^n;\mathbb{Z}/2) \cong K_*^M(k)\otimes\mathbb{Z}/2[\tau,y_1,...,y_n]\otimes\Lambda(x_1,...,x_n).$$

The above theorem also holds for $p=2$, considering the graded ring $grH^{*,*'}(B(\mathbb{Z}/2)^n;\mathbb{Z}/2)$.

We add another arguments for $p = 2$. Define $d_{n,i} \in \mathbb{Z}/2[x_1, ..., x_n]$ by

$$\Pi_{x \in \mathbb{Z}/2\{x_1,...,x_n\}}(t + x) = t^{2^n} + \sum t^{2^i} d_{n,i}$$

so that $d_{n,i}^2 = c_{n,i}$ and we have the isomorphism

$$\mathbb{Z}/2[x_1, ..., x_n]^{GL_n(\mathbb{Z}/2)} \cong \mathbb{Z}/2[d_{n,0}, ..., d_{n,n-1}].$$

Moreover for the regular real representation $reg_{\mathbb{R}} : (\mathbb{Z}/2)^n \to BO_{2^n}$, we have

$$\tau^{-2^{n-1}+2^{i-1}+1} reg_{\mathbb{R}}^*(w_{2^n-2^i}) = d_{n,i}$$

where we identify the element $d_{n,i}$ with a homogeneous element of $w(d_{n,i}) = 2$. We know from Lemma 5.7 in [Ka-Te-Ya]

$$d_{n,i} = Q_0...\hat{Q}_i...Q_{n-2}(x_1...x_n)$$

where $d_{n,i}$ is consider in $(*)$ (equivalently, in

$$H^{*,*'}((B\mathbb{Z}/2)^n; \mathbb{Z}/2)/(\tau \cdot H^{*-1,*'}(B(\mathbb{Z}/2)^n; \mathbb{Z}/2))$$

(see for details §5 in [Ka-Te-Ya]).

Theorem 5.2. *Let $a = a_1...a_n \in H^{n,n}(X; \mathbb{Z}/2)$ and $\tau a' = a$. Then we have*

$$\tau \cdot q_i(1)' = \tau \cdot Q_0...\hat{Q}_i...Q_{n-2}(a') = Q_0...\hat{Q}_i...Q_{n-2}(a)$$
$$= i_a^*(d_{n,i}) = i_a^*(\tau^{-2^{n-1}+2^{i-1}+1} reg_{\mathbb{R}}^*(w_{2^n-2^i})).$$

Remark. Note that when $a \in K_*^M(k)/2 \cong H^{*,*}(pt.; \mathbb{Z}/p)$, the value of the above equation is zero (since $Q_i a = 0$).

Remark. Smirnov and Vishik constructed the (generalized) Arason map $e_n : I^n/I^{n+1} \to K_*^M(k)/2$ where I is the fundamental ideal of the Witt ring $W^*(k)$ generated by even dimensional quadratic forms. They used the Nisnevich classifying space $BO(n)_{Nis}$. In this paper we write the etale classifying space by $BO(n)_{et}$ or simply $BO(n)$. The motivic cohomology is given as (Theorem 3.1.1 in [Sm-Vi])

$$H^{*,*'}(BO(n)_{Nis}; \mathbb{Z}/2) \cong H^{*,*'}(pt. : \mathbb{Z}/2) \otimes \mathbb{Z}/2[u_1, ..., u_n]$$

where $u_i = \tau^{-[(i+1)/2]} w_i$ identifying $w_i \in H^{i,i}(BO(n)_{et}; \mathbb{Z}/2)$. These elements are called as subtle characteristic classes in the paper [Sm-Vi].

Let X_q be the quadric for a quadratic form q, and $\chi_q = \chi_{X_q}$. Smirnov-Vishik proved (Theorem 3.2.34, Remark 3.2.35 in [Sm-Vi]) that when $q \in I^n$, there is a map $f_q : \chi_q \to BO(N)_{Nis}$ (with $N = dim(q)$), and the Arason map can be defined as (for $1 \leq i \leq n-2$)

$$e_n(q) = \tau(Q_0...\hat{Q}_i...Q_{n-2})^{-1} f_q^*(u_{2^n-2^i}).$$

We consider the following diagram

$$\begin{array}{ccccc} H^{*,*'}(\chi_q;\mathbb{Z}/2) & \xleftarrow{f_q^*} & H^{*,*'}(BO(N)_{Nis};\mathbb{Z}/2) & \xleftarrow{j^*} & H^{*,*'}(BO(N)_{et};\mathbb{Z}/2) \\ {\scriptstyle p^*}\downarrow & & & & \downarrow \\ H^{*,*'}(X_q:\mathbb{Z}/2) & \xleftarrow{i_a^*} & H^{*,*'}(B(\mathbb{Z}/2)^n;\mathbb{Z}/2) & \xleftarrow{reg^*} & H^{*,*'}(BO(2^n)_{et};\mathbb{Z}/2) \end{array}$$

where $N \geq max(dim(f), 2^n)$, and $p : X_q \to \chi_q$ and $j : BO(N)_{Nis} \to BO(N)_{et}$ are natural induced maps, and i_a^* is the induced map from $a = a_1...a_n \in H^n_{et}(X_q;\mathbb{Z}/2)$. Hence we have two elements

$$u = p^* f_q^*(u_{2^n-2^i}) = p^* f_q^*(\tau^{-2^{n-1}+2^{i-1}} \cdot j^*(w_{2^n-2^i}))$$

$$v = i^*_{e_n(q)}\tau^{-1}d_{n,i} = i^*_{e_n(q)}(\tau^{-2^{n-1}+2^{i-1}} \cdot reg^*_{\mathbb{R}}(w_{2^n-2^i})).$$

We see $\tau u = \tau v$, however this is worthless, since both sides are just zero, and the map i_a^* may lose information.

6. Lie groups G and the flag manifolds G/T

Let Y be a simply connected H-space of finite type (e.g. compact Lie group). By Borel, its mod p-cohomology is (for p odd)

$$H^*(Y;\mathbb{Z}/p) \cong P(y)/p \otimes \Lambda(x_1,...,x_\ell), \quad P(y) = \otimes_i^s \mathbb{Z}_{(p)}[y_i]/(y_i^{p^{r_i}})$$

where $|y_i|$ is even and $|x_j|$ is odd. When $p = 2$, a graded ring $grH^*(Y;\mathbb{Z}/2)$ is isomorphic to the right hand side ring, e.g. $x_j^2 = y_{i_j}$ for some y_{i_j}.

For ease of arguments, we assume p *odd* in this section (however the similar results also hold for p=2). Let us write by $QH^*(-)$ the indecomposable module

$$QH^*(Y;\mathbb{Z}/p) = H^*(Y;\mathbb{Z}/p)/(H^+(Y;\mathbb{Z}/p))^2 \cong \mathbb{Z}/p\{y_1,...,y_s,x_1,...,x_\ell\}.$$

Then it is known by Lin, Kane [Ka]

Lemma 6.1. *(J.Lin, §35 in [Ka]) Let p is an odd prime and Y be a finite simply connected H-space. Then*

$$Q^{even}H^*(X;\mathbb{Z}/p) = \sum_{n\geq 0} \beta P^n Q^{odd} H^*(X;\mathbb{Z}/p).$$

For connected but non simply connected space, the above theorem holds for $* \geq 4$. Moreover

Lemma 6.2. *(§36 in [Ka]) By the same assumption as the preceding lemma, we see*

$$Q^{2n}H^*(Y;\mathbb{Z}/p) = 0 \quad unless \quad n = p^k + ... + \hat{p}^i + ... + 1.$$

For example, the possibility of even degree of ring generators are

$$(p+1),\quad (p+1,p^2+1),\quad (p^2+p+1,p^3+p+1,p^3+p^2+1),...$$

Moreover, it is known that each generator is combined by reduced power operations.

The first degree type $|y| = 2(p+1)$ appears for $p = 3$, exceptional Lie groups F_4, E_6, E_7, and for $p = 5$, E_8 (and for $p = 2$, G_2, F_4, E_6). The second type appears for $p = 3$, E_8. However it seems that there is no example for other types for p odd.

So we assume the first or second types above. Let us say that they are of type (I) and (II) respectively. When type (I), there are x_1, x_2, y in $H^*(Y;\mathbb{Z}/p)$ such that

$$(*)\quad x_2 = P^1(x_1),\quad y = Q_1(x_1) = Q_0(x_2)$$

with $|x_1| = 3$, $|x_2| = 2p+1$ and $|y| = 2p+2$. The existence of the above y also holds for $p = 2$. For type (II), there are more elements x_3, x_4 and y' such that

$$(**)\quad y' = P^p(y) = Q_2(x_1) = Q_1(x_3) = Q_0(x_4),$$

where $|y'| = 2(p^2+1)$, $|x_3| = 2p^2-2p+3$, $|x_4| = 2p^2+1$. For $p = 2$, the other types appear but we see ;

Lemma 6.3. *Let G be a simply connected Lie group such that $H^*(G;\mathbb{Z})$ has p-torsion. Then there are x_1, x_2, y satisfying $(*)$.*

Let T be the maximal torus of a simply connected compact Lie group G and BT the classifying space of T. We consider the fibering

$$(6.1)\quad G \xrightarrow{\pi} G/T \xrightarrow{i} BT$$

and the induced spectral sequence

$$E_2^{*,*} = H^*(BT; H^*(G;\mathbb{Z}/p)) \Longrightarrow H^*(G/T;\mathbb{Z}/p).$$

The cohomology of the classifying space of the torus is given by

$$H^*(BT) \cong S(t) = \mathbb{Z}[t_1,...,t_\ell] \quad with \quad |t_i| = 2.$$

where ℓ is also the number of the odd degree generators x_i in $H^*(G;\mathbb{Z}/p)$. It is known that y_i are permanent cycles and that there is a regular sequence ([Tod],[Mi-Ni]) $(\bar{b}_1,...,\bar{b}_\ell)$ in $H^*(BT)/(p)$ such that $d_{|x_i|+1}(x_i) = \bar{b}_i$. Thus we get

$$E_\infty^{*,*'} \cong grH^*(G/T;\mathbb{Z}/p) \cong P(y)/p \otimes S(t)/(\bar{b}_1,...,\bar{b}_\ell).$$

Moreover we know that G/T is a manifold of torsion free, and

$$(6.2)\quad H^*(G/T)_{(p)} \cong \mathbb{Z}_{(p)}[y_1,..,y_k] \otimes S(t)/(f_1,...,f_k,b_1,...,b_\ell)$$

where $b_i = \bar{b}_i \ mod(p)$ and $f_i = y_i^{p^{r_i}} \ mod(t_1, ..., t_\ell)$. Since $H^*(G/T)$ is torsion free, we also know

$$BP^*(G/T) \cong BP^*[y_1, ..., y_k] \otimes S(t)/(\tilde{f}_1, ..., \tilde{f}_k, \tilde{b}_1, ..., \tilde{b}_\ell) \tag{6.3}$$

where $\tilde{b}_i = b_i \ mod(BP^{<0})$ and $\tilde{f}_i = f_i \ mod(BP^{<0})$.

Let G_k be the split reductive algebraic group corresponding G and T_k the split maximal torus. Hence $CH^*(G_k) \cong CH^*(G_{\mathbb{C}})$. We also have $CH^*(G_k/T_k) \cong H^*(G/T)$ and $CH^*(BT_k) \cong H^*(BT)$.

Next we consider the relation between $CH^*(G_k)$ and $CH^*(G_k/T_k)$.

Theorem 6.4. *(Grothendieck [Gr]) Let $\mathbb{G}_k$ be a G_k-torsor over k. (Here we do not assume the non triviality of $\mathbb{G}_k$). Let $h^*(-)$ be an oriented cohomology (e.g., $CH^*(X), \Omega^*(X)$). Then*

$$h^*(\mathbb{G}_k) \cong h^*(\mathbb{G}_k/T_k)/(i^*h^*(BT_k)) \cong h^*(\mathbb{G}_k/T_k)/(t_1, ..., t_\ell).$$

Since $CH^*(G_k) \cong CH^*(G_{\mathbb{C}})$, from the result $H^*(G/T)$ (6.2), we have

Corollary 6.5. *(Kac [Ka]) $CH^*(G_k)_{(p)} \cong P(y)/(py_i)$.*

Proof for groups of type (I) or (II). From the above theorem, we only need to show $py = py' = 0$. By the integral Beilinson-Lichtembaum conjecture (solved by Voevodsky), we can take $x_1 \in H^{3,2}(G_k; \mathbb{Z}/p)$ (note $x_1 \in H^3(G; \mathbb{Z})$) so that

$$Q_1(x) = y \in H^{2*,*}(G_k; \mathbb{Z}/p) = CH^*(G_k)/p,$$

That is $y = Q_0P^1(x_1)$ and y is p-torsion also in $CH^*(G_k)$. The element $y' = Q_0P^pP^1(x_1)$ is also p-torsion for type (II) groups. □

Corollary 6.6. *If $CH^*(\mathbb{G}_k/T_k)$ is generated as a ring by t_i, then $CH^*(\mathbb{G}_k) \cong \mathbb{Z}_{(p)}$.*

It is known that we can take generators such that $Q_i(x_{odd}) \in P(y_{even})/(p)$ for all $i \geq 0$ ([Mi-Ni]). The following result for $\Omega^*(G_{\mathbb{C}})$ is one of the main results in [Ya1] which is proved by using Lemma 4.4.

Theorem 6.7. *([Ya1]) Take generators so that $Q_i(x_{odd}) \in P(y_{even})/(p)$ for all $i \geq 0$. Then there is an Ω^*-module isomorphism*

$$\Omega^*(G_k)/I_\infty^2 \cong \Omega^* \otimes P(y_{even})/(I_\infty^2, \sum_i v_iQ_i(x_{odd})).$$

Corollary 6.8. *For a simply connected simple group G of type (I), we have $\Omega^*(G_k) \cong \Omega^*[y]/(py, v_1y, y^p)$. For type (II), we have*

$$\Omega^*(G_k)/I_\infty^2 \cong \Omega^*[y, y']/(I_\infty^2, py, v_1y + v_2y', py', v_1y', y^p, (y')^p).$$

Corollary 6.9. *When $X = G_k$ of type (I) or (II), we have*

$$tor_v(y) = tor_v(y') = 2 \quad but \quad tor_v((y')^2) = 3.$$

Proof. For type (II), we have the relation $v_1 y + v_2 y' = 0$ in $\Omega^*(X)$. This means that $v_1 y = 0 \in gr\Omega^*(X)$ since $|y| < |y'|$, while $v_2 y' \neq 0$ in $gr\Omega^*(X)$. We have $v_2(y')^2 = -v_1 y y' = 0$, since $v_1 y' = 0$. □

7. twisted flag variety $\mathbb{G}_k/T_k$

Recall that $\mathbb{G}_k$ is the nontrivial G_k-torsor. We can construct the twisted form of G_k/T_k by

$$(\mathbb{G}_k \times G_k/T_k)/G_k \cong \mathbb{G}_k/T_k.$$

We will study the twisted flag variety $X = \mathbb{G}_k/T_k$.

Let $dim(G/T) = 2d$. Then the torsion index is defined as

$$t(G) = H^{2d}(G/T; \mathbb{Z})/H^{2d}(BT; \mathbb{Z}).$$

Let $n(\mathbb{G}_k)$ be the greatest common divisor of the degrees of all finite field extension k' of k such that $\mathbb{G}_k$ becomes trivial over k'. Then by Grothendieck, it is known that $n(\mathbb{G}_k)$ divides $t(G)$. Moreover, there is a G_k-torsor $\mathbb{G}_F$ over some extension field F of k such that $n(\mathbb{G}_F) = t(G)$. Note that $t(G_1 \times G_2) = t(G_1) \cdot t(G_2)$. It is well known that if $H^*(G)$ has a p-torsion, then p divides the torsion index $t(G)$. Torsion index for simply connected compact Lie groups are completely determined by Totaro [To1,2]. Hence we also see that there is k such that there is non trivial G_k-torsor $\mathbb{G}_k$. In particular, we have

Lemma 7.1. *Let G be a group of type (I). Then $t(G)_{(p)} = p$.*

To seek the condition for $v_n y \in Res_\Omega$, we recall the (integral) Morava K-theory $\tilde{K}(n)^*(-)$ with the coefficient

$$\tilde{K}(n)^*(pt.) = \tilde{K}(n)^* = \mathbb{Z}_{(p)}[v_n, v_n^{-1}].$$

We also consider its $\mathbb{A}^1$-homotopy version

$$(*) \quad A\tilde{K}(n)^{2*,*}(X) \cong \Omega^*(X) \otimes_{BP^*} \mathbb{Z}_{(p)}[v_n, v_n^{-1}].$$

Lemma 7.2. *Let y be an element with (1),(2) in §3 such that $y \notin Res_\Omega$ but $y \in Res_{\tilde{K}(n)}$. Moreover assume $|y| \leq 4(p^n - 1)$. Then $deg_v(y) = n + 1$.*

Proof. Each Ω^*-module generator of $\Omega^*(\bar{X})$ is also a $\tilde{K}(n)^*$-module generator of $A\tilde{K}(n)^{2*,*}(\bar{X})$. By the assumption, there is $y' \in A\tilde{K}^{2*,*}(X)$ with $Res_{\tilde{K}(n)}(y') = y$ from $(*)$. This means that there is $y_\Omega \in \Omega^*(X)$

such that $v_n^s y_\Omega = y'$ for $s \in \mathbb{Z}$. Since $y = res_\Omega(v_n^s y_\Omega)$, the assumption $y \notin Res_\Omega$ implies $s < 0$. From the dimensional condition, we see

$$0 < |y_\Omega| = |y'| + 2s(p^n - 1) \leq (2s+4)(p^n - 1).$$

The only case is $s = -1$ and we get $y_\Omega = v_n y$, i.e., $deg_v(y) \geq n+1$. □

Here we note for the Morava K-theory of the Rost motive R_n.

Lemma 7.3. *Let $R_a = R_n$, and $0 \leq m \leq n-1$. Then the restriction map $res_{\tilde{K}(m)}$ is an isomorphism*

$$A\tilde{K}(m)^{2*,*}(R_a) \cong A\tilde{K}(m)^{2*,*}(\bar{R}_a).$$

Proof. From Theorem 4.6, we have

$$CH^*(R_n) \cong \mathbb{Z}_p\{1\} \oplus M_n(y)[y]/(y^{p-1}).$$

Recall $\tilde{K}(m)^* = \mathbb{Z}_{(p)}[v_m, v_m^{-1}]$. Then we have

$$A\tilde{K}(m)^{2*,*}(R_a) \cong \tilde{K}(m)^{2*} \oplus \tilde{K}(m)^{2*}\{py, v_1 y, ..., v_{n-1}y\}[y]/(y^{p-1})$$

$$\cong \tilde{K}(m)^{2*} \oplus \tilde{K}(m)^{2*}\{py, v_m y\}[y]/(y^{p-1}) \cong \tilde{K}(m)^{2*} \oplus \tilde{K}(m)^{2*}\{v_m y\}[y]/(y^{p-1})$$

$$\cong \tilde{K}(m)^{2*}[y]/(y^p) \cong \tilde{A}K(m)^{2*,*}(\bar{R}_a).$$

□

Recall the usual Morava k-theory with the coefficients $K(m)^* \cong \mathbb{Z}/p[v_m, v_m^{-1}]$. Then $AK(m)^{2*,*}(R_n) \cong A\tilde{K}(m)^{2*,*}(R_2)/p$ and we also have the isomorphism $AK(m)^{2*,*}(R_a) \cong AK(m)^{2*,*}(\bar{R}_a)$.

In particular, $res_{\tilde{K}(1)}$ is always surjective for $X = R_2$. Similar fact holds for $X = \mathbb{G}_k/T_k$.

Lemma 7.4. *For a simply connected Lie group G having p-torsion in cohomology, the following restriction map is surjective ;*

$$A\tilde{K}(1)^{2*,*}(\mathbb{G}_k/T_k) \to A\tilde{K}(1)^{2*,*}(G_k/T_k) \to \tilde{K}(1)^*\{y\}.$$

Proof. We prove the case of Type (I) and the other types are proved similarly. Since we have the classifying map $\mathbb{G}_k/T_k \to BT_k$, each element t_i from $h^*(BT_k)$ is in the image Res_h for all orientable cohomology $h^*(-)$. In particular, $t_i \in Res_{\tilde{K}(1)}$. From Corollary 6.8, $v_1 y = 0 \in \Omega^*(G_k)$ and this implies $v_1 y$ is written by elements $\sum v_1^r t_{i_1}...t_{i_s}$ from Theorem 6.4. Hence $v_1 y \in Res_{\tilde{K}(1)}$. □

Remark. More strongly, it is well known by Hodgikin [Ho] that $K^0_{top}(G/T)$ (and hence $\tilde{K}(1)^*(G/T)$) is multiplicatively generated by t_i (and so the image from $K^0_{top}(BT)$).

Of course $|y| = 2(p+1) < 4(p^2-1)$, from Lemma 7.2, we get

Corollary 7.5. *Let G be a simply connected semisimple Lie group having p-torsion in cohomology. Then there is a nontrivial G_k torsor $\mathbb{G}_k$ and there is $0 \neq y \in CH^{2p+2}(G_k/T_k)$ such that $deg_v(y) = 2$, so that*

$$\mathbb{Z}_{(p)}\{c_0\} \oplus \mathbb{Z}/p\{c_1\} \twoheadleftarrow CH^*(M) \twoheadleftarrow CH^*(\mathbb{G}_k/T_k)$$

for some irreducible motive M.

For groups of type (I), we show more strong results in §9, 10 below.
Remark. Garibaldi and Zainoulline show also more strong result. For $p = 3$ or 5, the p-torsion part is isomorphic to $\mathbb{Z}/p$ (Proposition 5.4 in [Ga-Za]) for simply connected simple group G. This follows from their result (Theorem in [Ga-Za]) : the torsion part of the graded ring for the γ-filtration of K-theory (see §15 below)

$$torsion(gr^4_\gamma(\mathbb{G}_k/T_k)) \cong \mathbb{Z}/N(G)$$

where $N(G)$ is the Dynkin index of G. For example, when $G = E_8$, the Dynkin index $N(G) = 60$.

The Q_i-action on the motivic cohomology $H^{*,*'}(\mathbb{G}_k/T_k; \mathbb{Z}/p)$ seems to be similar to that on the norm variety (or the Rost motive R_2).

Lemma 7.6. *Let G be a Lie group with p-torsion in cohomology and $X = \mathbb{G}_k/T_k$. Then there is a field extension k'/k, a pure symbol $a \in K^M_3(k')/p$ and $a' \in H^{3,2}(X; \mathbb{Z}/p)$ with $\tau a' = a$ such that in the motivic cohomology $H^{*,*'}(X; \mathbb{Z}/p)$ over k', we have*

$$Q_1(a') = c_0(y), \quad Q_0(a') = c_1(y).$$

Proof. In $CH^*(X)$, we see $pc_1(y) = 0$, and so in $\Omega^*(X)/BP^{<0}$, namely there is a relation

$$pc_1(y) + v_1c' = 0 \quad in\ \Omega^*(X)/(I^2_\infty, v_2, v_3, ...).$$

Then we see $v_1res(c') = pv_1y \in \Omega^*(\bar{X})/(I^3_\infty, v_2, ...)$ and so $res(c') = py \in \Omega^*(X)/(I^2_\infty, v_2, ...)$. Hence we can take c' as $c_0(y)$. Then from Lemma 4.4, there is $b \in H^{3,2}(X; \mathbb{Z}/p)$ such that

$$Q_1(b) = c' = c_0(y), \quad and \quad Q_0(b) = c_1(y).$$

Let $a = \tau b \in H^{3,3}(X; \mathbb{Z}/p)$. Then $a \neq 0$ since it is well known ([Or-Vi-Vo], [Vo]) that $H^{n+1,n}(Y; \mathbb{Z}/p) \xrightarrow{\times\tau} H^{n+1,n+1}(Y; \mathbb{Z}/p)$ is injective for each smooth Y.

Consider the well known spectral sequence for the etale cohomology

$$E_2^{*,*'} \cong H^*_{et}(Spec(k); H^*(G_k/T_k; \mathbb{Z}/p)) \Longrightarrow H_{et}(\mathbb{G}_k/T_k; \mathbb{Z}/p).$$

We see that $a \in H^{3,3}(X; \mathbb{Z}/p) \cong H^3_{et}(X; \mathbb{Z}/p)$ can be represented as an element in the sub-quotient ring of E_2-term, which is generated as a

ring by $H^*(G/T)$ and $H^*_{et}(pt;\mathbb{Z}/p)$. Hence we can write

$$a = \sum \tau t_i a'_i + \sum a_1 a_2 a_3$$

for $a'_i, a_i \in K^M_1(k)/p \cong H^1_{et}(Spec(k);\mathbb{Z}/p)$. Here note $t_i a'_i \in H^{3,2}(X;\mathbb{Z}/p)$ and $Q_i(t_i a'_i) = 0$ since the Q_i operations on both t_i and a'_i are zero. Taking off $a_i t_i$ from b, we can take

$$a = \sum a_1 a_2 a_3 \in K^M_3(k)/p.$$

It is known from Theorem 2.10 in [Or-Vi-Ve] or Corollary 8.4 in [Ya6] that we can take a field k' such that $a|_{k'}$ becomes a pure symbol. This shows the lemma. □

8. generically splitting variety

Let $P \supset T$ be a parabolic subgroup of G. Petrov, Semenov and Zainoulline developed the theory of decompositions of motives $M(\mathbb{G}_k/P_k)$. In this section, we recall their results.

By extending the arguments by Vishik in quadrics to flag varieties, Petrov, Semenov and Zainoulline define the J-invariant of $\mathbb{G}_k$. Recall the expression in §6

$$(*) \quad H^*(G;\mathbb{Z}/p) \cong \mathbb{Z}/p[y_1,...,y_s]/(y_1^{p^{r_1}},...,y_s^{p^{r_s}}) \otimes \Lambda(x_1,...,x_\ell).$$

The J-invariant is defined as

$$J_p(\mathbb{G}_k) = (j_1,...,j_s)$$

if j_i is the minimal integer such that

$$y_i^{p^{j_i}} \in Im(res_{CH}) \quad mod(y_1,...,y_{i-1},t_1,...,t_\ell)$$

for $res_{CH} : CH^*(\mathbb{G}_k/T_k) \to CH^*(G_k/T_k)$. Here we take $|y_1| \le |y_2| \le ...$ in $(*)$. Hence $0 \le j_i \le r_i$ and $J_p(\mathbb{G}_k) = (0,...,0)$ if and only if $\mathbb{G}_k$ split by an extension of the index coprime to p. One of the main results in [Pe-Se-Za] is

Theorem 8.1. *(Theorem 5.13 in [Pe-Se-Za]) Let $\mathbb{G}_k$ be a G_k-torsor over k and $J_p(\mathbb{G}_k) = (j_1,...,j_s)$. Then there is a mod$(p)$ motive $R(\mathbb{G}_k;\mathbb{Z}/p)$ such that (additively)*

$$CH^*(R(\mathbb{G}_k;\mathbb{Z}/p)|_{\bar{k}})/p \cong \mathbb{Z}/p[y_1,...,y_s]/(y_1^{p^{j_1}},...,y_s^{p^{j_s}}),$$

$$M(\mathbb{G}_k/B_k;\mathbb{Z}/p) \cong \oplus_u R_p(\mathbb{G}_k;\mathbb{Z}/p) \otimes \mathbb{T}^{\otimes u}$$

$$\cong R(\mathbb{G}_k;\mathbb{Z}/p) \otimes \mathbb{Z}/p[y_1^{p^{j_1}},...,y_s^{p^{j_s}}]/(y_1^{p^{r_1}},...,y_s^{p^{r_s}}) \otimes S(t)/(p,\bar{b}_1,...,\bar{b}_\ell)).$$

Note that G is of type I means that

$$r = 1, \quad d = |y|/2 > 1 \quad (\textit{and hence } k = 1)$$

in the notation (see §7.2 for details) in [Pe-Se-Za].

Corollary 8.2. *Let G be a group of type (I). Then $J(\mathbb{G}_k) = (1)$ and*

$$CH^*(\bar{R}(\mathbb{G}_k; Z/p))/p \cong \mathbb{Z}/p[y]/(y),$$
$$M(\mathbb{G}_k/T_k; \mathbb{Z}/p) \cong \oplus_u R(\mathbb{G}; \mathbb{Z}/p) \otimes \mathbb{T}^{\otimes u}$$

where we can identify the sum of the Tate motive

$$\oplus_u \mathbb{T}^{\otimes u} \cong H^*(G/T; \mathbb{Z}/p)/(y) \cong S(t)/(p, \bar{b}_1, ..., \bar{b}_\ell).$$

Corollary 8.3. *For a group G of type (I), we have*

$$\mathbb{Z}/p\{1\} \oplus M_2(y)/p \cong \mathbb{Z}/p\{1, c_0, c_1\} \twoheadleftarrow R(\mathbb{G}; \mathbb{Z}/p),$$

and $deg(yt) = 2$ for each $0 \neq t \in CH^*(\bar{X})/(y)$.

Petrov, Semenov and Zainoulline develop the theory of generically splitting varieties. We say that L is splitting field of a variety of X if $M(X|_L)$ is isomorphic to a direct sum of twisted Tate motives $\mathbb{T}^{\otimes i}$ and the restriction map $i_L : M(X) \to M(X|_L)$ is isomorphic after tensoring $\mathbb{Q}$. A smooth scheme X is said to be generically split over k if its function field $L = k(X)$ is a splitting field. Note that (the complete flag) $X = \mathbb{G}_k/B_k$ is always generically split.

Theorem 8.4. *(Theorem 3.7 in [Pe-Se-Za]) Let $Q_k \subset P_k$ be parabolic subgroups of G_k which are generically split over k. Then there is a decomposition of motive : $M(\mathbb{G}/Q_k) \cong M(\mathbb{G}_k/P_k) \otimes H^*(P/Q)$.*

In particular, when G is of type (I) (i.e., $r = 1$ and $d > 1$ in the notation [Pe-Se-Za]), from §7 in [Pe-Se-Za], we have the *integral* (localized at p) motivic decomposition which deduces the $mod(p)$ decomposition in Theorem 7.1. Moreover we know (Corollary 6 in [Vi-Za], or §7 in [Pe-Se-Za]) that the integral motive corresponding $R(\mathbb{G}_k)_p$ is really the Rost motive R_2 localized at p.

9. Type (I) groups

We first consider the exceptional Lie groups

$$(*) \quad G = G_2, F_4, E_8 \quad for \quad p = 2, 3, 5 \ respectively.$$

The following fact is well known

Lemma 9.1. *([Mi-To]) Let G be a group in $(*)$. Then we have*

$$H^*(G; \mathbb{Z}/p) \cong \mathbb{Z}/p[y]/(y^p) \otimes \Lambda(x_1, ..., x_\ell)$$

with $\ell = 2(p-1)$ and $P^1 x_{2i-1} = x_{2i}$ for $1 \leq i \leq p-1$.

Proof. It is known that for a simply connected simple exceptional Lie group if there are x_i, $x_{i'}$ with $deg(x_i) - deg(x_{i'}) = 2(p-1)$, then we can take generators such that $P^1(x_{i'}) = x_i$ ([Mi-To]). For the group in $(*)$, odd degree generator x is written $x_1, ..., x_{2(p-1)}$ such that $|x_{2i}| = |y^i| - 1 = 2(p+1)i - 1$ and $|x_{2i}| - |x_{2i-1}| = 2(p-1)$.

For example $H^*(E_8; \mathbb{Z}/5)$ is isomorphic to

$$\mathbb{Z}/5[y_{12}]/(y_{12}^5) \otimes \Lambda(x_3, x_{11}, x_{15}, x_{23}, x_{27}, x_{35}, x_{39}, x_{47})$$

where suffix means its degree so that

$$P^1 x_3 = x_{11}, \quad P^1 x_{15} = x_{23}, \quad P^1 x_{27} = x_{35}, \quad P^1 x_{39} = x_{47}.$$

□

Since $py^i = 0$ in $H^*(G; \mathbb{Z}_{(p)})$, we know $py^i = 0$ $mod(Ideal(S(t))$ in $H^*(G/T; \mathbb{Z}_{(p)})$. We need the following more strong result

$$py^i = b_{2i} \in S(t) \cong H^*(BT; \mathbb{Z}).$$

Recall that $\bar{b}_j$ is an element in $S(t)/p$ so that $d_{|x_j|+1}(x_j) = \bar{b}_j$ in the Serre spectral sequence converging to $H^*(G/T; \mathbb{Z}/p)$. Hence $b_{2i} = \bar{b}_{2i}$ $mod(p)$. The above assumption holds when $p = 2, 3$ from the direct observations (see §10 below). For $E_8, p = 5$, it seems not so easy to this property, while Nakagawa [Na] computed the integral cohomology completely. However after written the first version of this paper, Karpenko proved the following strong result

Theorem 9.2. *(Karpenko Lemma 2.1 in [Ka]) Let P_k be a parabolic subgroup of G_k and $\mathbb{G}_k/P_k$ be a generically twisted flag variety. Then the natural map $CH^*(BP_k) \to CH^*(\mathbb{G}_k/P_k)$ is surjective.*

Corollary 9.3. *For group G in $(*)$, the Chow ring $CH^*(\mathbb{G}_k/B_k)$ is generated by elements t_i in $S(t)$. In particular, $py^i \in S(t)$.*

Proof. For some extension F/k, of order ap, a is coprime p, the G_k-torsor $\mathbb{G}_k$ splits. Hence $py^i \in Im(res_{CH} : CH^*(\mathbb{G}_k/B_k) \to CH^*(G_k/B_k))$, which is written by elements in $S(t)$ by the above Karpenko theorem □

Since $P^1(x_{2i-1}) = x_{2i}$ from Lemma 9.1, we have

$$P^1(\bar{b}_{2i-1}) = \bar{b}_{2i} \neq 0 \in H^*(BT; \mathbb{Z}/p)$$

by the transgression theorem. So there is $b_{2i-1} \in BP^*(BT)$ such that

$$r_{\Delta_1}(b_{2i-1}) = b_{2i} \neq 0 \in BP^*(BT)/I_\infty.$$

Since $py^i = b_{2i}$ in $H^*(G/T)$, the above fact implies

$$r_{\Delta_1}(b_{2i-1}) = py^i \neq 0 \in BP^*(G/T)/(I_\infty BP^*(BT)).$$

Lemma 9.4. *We have*

$$b_{2i-1} = v_1(y^i + t) \in BP^*(G/T)/(I_\infty^2, v_2, v_3, ...) \quad for\ t \in S(t).$$

Hence $deg_v(y^i) = 2$ *and we can take* $c_1(y^i) = b_{2i-1}$.

Proof. Recall $grBP^*(G/T) \cong BP^*[y, t_1, ..., t_\ell]/(f_1, \tilde{b}_1, ..., \tilde{b}_2)$ where $f_1 = y^p\ mod(I_\infty, S(t))$ and $\tilde{b}_i = \bar{b}_i\ mod(I_\infty)$. Since $b_{2i-1} = 0$ in $H^*(G/T; \mathbb{Z}/p)$ we can write

$$b_{2i-1} = \sum_{j<i}(py^j t(j) + v_1 y^j t(j)') + \lambda v_1 y^i + v_1 t \quad in\ (I_\infty^2, v_2, ...)BP^*(G/T)$$

where $\lambda \in \mathbb{Z}/p$, and $t(j), t(j)', t \in S(t)$. We will prove $\lambda \neq 0\ mod(p)$.

By induction on i, we can assume $v_1 y^j = b_{2j-1}$ for $j < i$. Note $r_{\Delta_1}(y) = 0\ mod(I_\infty)$. By the Cartan formula of the operation r_{Δ_1}, we have

$$r_{\Delta_1}(b_{2i-1}) = \sum_{j<i}(py^j t(j)'') + \lambda p y^i + pt \quad in\ (p^2, I_\infty^{<0})BP^*(G/T).$$

Here $r_{\Delta_1}(b_{2i-1}) = b_{2i} = py^i$. Since $BP^*(G/T)/I_\infty^{<0} \cong H^*(G/T)$ is p-torsion free, we get

$$y^i = \sum_{j<i} y^j t(j)'' + \lambda y^i + t$$

in $H^*(G/T)/p \cong H^*(G/T; \mathbb{Z}/p)$. Hence $\lambda = 1\ mod(p)$, otherwise there is a relation $y^i = \sum_{j<i} y^j t(j)'' + t$ in $H^*(G/T; \mathbb{Z}/p)$, which is a contradiction to (6.2).

Thus we have the first equation in the lemma. The second equation follows from $b_{2i-1} \in BP^* \otimes S(t)$ and the dimensional reason $|y^i| \leq -|v_2| = 2(p^2 - 1)$, which shows $deg_v(y^i) \leq 2$. □

Remark. By Petrov, Semov and Zainoulline, the motive $M(\mathbb{G}_k/T_k)$ is isomorphic to a sum of the Rost motive R_2. Hence $deg_v(y^i) = 2$ is almost immediate.

Theorem 9.5. *Let* (G, p) *be the type* $(*)$. *Then* $c_i(y^j) = b_{2j-i} \in S(t)$ *for* $0 \leq i \leq 1 \leq j \leq p-1$ *and there is a ring isomorphism*

$$CH^*(\mathbb{G}_k/T_k)/p \cong S(t)/(b_j b_{j'} | 1 \leq j, j' \leq p-1)$$
$$\cong S(t)/(p, c_i(y^j)c_{i'}(y^{j'}) | 0 \leq i, i' \leq 1 \leq j, j' \leq p-1).$$

Proof. Let us write

$$\bar{S}(t) = H^*(G/T; \mathbb{Z}/p)/(y) \cong S(t)/(p, \bar{b}_1, ..., \bar{b}_{2p-2}).$$

From Theorem 7.1 (by Prtrov, Semenov, Zainoulline), we know

$$CH^*(\mathbb{G}_k/T_k)/p \cong CH^*(R(\mathbb{G}_k; \mathbb{Z}/p)) \otimes H^*(G/T; \mathbb{Z}/p)/(y)$$

$$\cong (\mathbb{Z}/p\{1\} \oplus M_2[y]/(y^{p-1})) \otimes \bar{S}(t)$$
$$\cong (\mathbb{Z}/p\{1, c_0(y), c_1(y), ..., c_0(y^{p-1}), c_1(y^{p-1})\}) \otimes \bar{S}(t).$$

On the other hand, we have an additive isomorphism

$$S(t)/(p, b_i b_j) \cong (\mathbb{Z}/p\{1, b_1, ..., b_{2p-2}\}) \otimes \bar{S}(t).$$

Here we see $c_1(y^i) = b_{2i-1}$ from Lemma 9.4, and $c_0(y^i) = b_{2i}$ from Karpenko' result Corollary 9.3. Therefore we have the isomorphism in this theorem by $c_0(y^i) \mapsto b_{2i}$ and $c_1(y) \mapsto b_{2i-1}$.

Thus we have the additive map, and it induces the ring map by the following reason. For example, the relation $c_i(y^j)c_{i'}(y^{j'}) = 0$ follows from $v_i y^j v_{i'} y^{j'} = 0 \ mod(I_\infty^2)$. □

Corollary 9.6. *Let G be the group of $(*)$. Then there is an isomorphism*

$$\Omega^*(\mathbb{G}_k/T_k)/(I_\infty^2) \cong \Omega^* \otimes S(t)/(Rel., I_\infty^2)$$

where the relation $(Rel.)$ is generated by

$$b_{2i}b_{2j} = pb_{2(i+j)}, \quad b_{2i-1}b_{2j-1} = v_1 b_{2(i+j)},$$

$$v_1 b_1 = pb_2, \quad b_{2i-1}b_{2j} = v_1 b_{2(i+j)} = pb_{2(i+j)-1}$$

letting $b_k = 0$ for $k \geq 2p-1$.

Proof. By the result by Petrov, Semenov, Zainoulline and Theorem 4.7, the restriction map

$$res_\Omega : \Omega^*(\mathbb{G}_k/T_k) \to \Omega^*(G_k/T_k)$$

is injective. Its image is

$$\Omega^*\{1, v_1 y, py, ..., v_1 y^{p-1}, py^{p-1}\} \otimes S(t)/(\tilde{b}_1, ..., \tilde{b}_{2p-2})$$
$$\cong \Omega^*\{1, b_1, ..., b_{2p-2}\}/(pb_{2i-1} - v_1 b_{2i-1}) \otimes S(t)/(\tilde{b}_1, ..., \tilde{b}_{2p-2}) \quad (1)$$

We see that the first relation holds in $\Omega^*(G_k/T_k)$ by

$$b_{2i}b_{2j} = py^i py^j = p^2 y^{i+j} = pb_{2(i+j)}.$$

The other relations are showed similarly. Hence we have a map

$$\Omega^* \otimes S(t)/(Rel., I_\infty^2) \to \Omega^*(\mathbb{G}_k/T_k)/(I_\infty^2).$$

Since there are relations $b_i b_j = ...$ and $pb_{2i-1} - v_1 b_{2i}$ in Rel, we have a map

$$f : (1) \to \Omega^* \otimes S(t)/(Rel., I_\infty^2).$$

The composition $res_\Omega \cdot f$ is an isomorphism, and we have the corollary. □

Given $p = 2, 3$ or 5, let $G(*)$ be the group in $(*)$ (e.g. $G(*) = F_4$ for $p = 3$). Let G be a simply connected simple Lie group of type (I). Then there is an embedding $j : G(*) \subset G$ such that its mod p cohomology has the same polynomial parts i.e., $P(y)/p \cong \mathbb{Z}/p[y]/(y^p)$ and moreover we have the ring isomorphism

$$H^*(G; \mathbb{Z}/p) \cong H^*(G(*); \mathbb{Z}/p) \otimes \Lambda(x_{2p-1}, ..., x_\ell).$$

Hence for $S(t') \cong \mathbb{Z}/p[t_{2p-1}, ..., t_\ell]$, we have

$$grH^*(G/T; \mathbb{Z}/p) \cong (H^*(G(*)/T(*); \mathbb{Z}/p) \otimes S(t'))/(\bar{b}_{2p-1}, ..., \bar{b}_\ell).$$

From the result (Theorem 7.1) by Petrov, Semenov and Zainoulline, we have the map $j^* : R(G; \mathbb{Z}/p) \to R(G(*); \mathbb{Z}/p)$ which induces the isomorphism

$$\bar{R}(G; \mathbb{Z}/p) \cong \bar{R}(G(*); \mathbb{Z}/p) \cong \mathbb{Z}/p[y]/(y^p).$$

We can show that j^* is an isomorphism. To see this, we recall the following theorem by Vishik and Zainoulline (Theorem 1 in [Vi-Za]) ; Let $f : M \to N$ be a map of motives with N in $M(X)$. If f is split surjective on $k(X)$, then so is f itself.

Hence the above map j^* is split surjective. So it is an isomorphism since $R(G; \mathbb{Z}/p)$ is an irreducible motive, namely

$$R(G; \mathbb{Z}/p) \cong R(G(*); \mathbb{Z}/p).$$

Therefore we have

Theorem 9.7. *Let G be a simply connected simple Lie group. Let $X = \mathbb{G}_k/T_k$ for nontrivial torsor $\mathbb{G}_k$. Then*

$$grCH^*(X)/p \cong (CH^*(G(*))/p \otimes S(t'))/(\bar{b}_{2p-1}, ..., \bar{b}_\ell)$$
$$\cong S(t)/(p, \bar{b}_i\bar{b}_j, \bar{b}_k | 1 \leq i, j \leq 2p-2 < k \leq \ell).$$

From Corollary 6.6, we have :

Corollary 9.8. *Let G be a group in the above theorem. Then we have the isomorphism $CH^*(\mathbb{G}_k)/p \cong \mathbb{Z}/p$.*

Let $gr_\gamma(X)$ be the graded ring associated by the γ-filtration of $K^0(\bar{X})$. (See §15 below for γ-filtration.)

Corollary 9.9. *Let G be a group in the above theorem. Then we have the isomorphism $gr_\gamma(\bar{X})/p \cong CH^*(X)/p$.*

We note that the above two corollaries are just a corollary of Karpenko's theorem (Theorem 9.2). These holds for all generically twisted flag variety $\mathbb{G}_k/B_k$ over k.

10. $G_2, p = 2$ AND $F_4, p = 3$

In this section, we write down each b_j for the groups G_2, F_4. We consider the integral ($\mathbb{Z}_{(p)}$) cohomology. We first study the case $(G, p) = (G_2, 2)$. We recall the cohomology from Toda-Watanabe [To-Wa]

$$H^*(G/T;\mathbb{Z}) \cong \mathbb{Z}[t_1, t_2, y]/(u, t_2^3 - 2y, y^2) \quad u = t_1^2 + t_1t_2 + t_2^2$$

with $|t_i| = 2$ and $|y| = 6$. The following theorem is just a corollary of Theorem 9.5 in the preceding section, however we write down the direct proof, it is get more easily.

Theorem 10.1. *([Ya5]) We have the ring isomorphisms*

$$CH^*(\mathbb{G}_k/T_k)_{(2)} \cong \mathbb{Z}_{(2)}[t_1, t_2]/(t_2^6, 2u, t_2^3u, u^2)$$

$$\cong \mathbb{Z}_{(2)}[t_1, t_2]/(c_0^2, 2c_1, c_0c_1, c_1^2)$$

identifying $u = t_1^2 + t_1t_2 + t_2^2 = v_1y = c_1$ *and* $c_0 = 2y = t_2^3$.

Proof. Since $u = 0$ and $t_2^3 - 2y = 0$ in $H^*(\bar{X})$, we can take $c_1 = u$ (which is represented by a Chern class) and $c_0 = t_2^3$ in $CH^*(X)$. Additively we can write

$$(*) \quad CH^*(X) \cong (\mathbb{Z}\{1, c_0\} \oplus \mathbb{Z}/2\{c_1\}) \otimes H^*(G/T;\mathbb{Z})/(y)$$

$$\cong (\mathbb{Z}\{1, t_2^3\} \oplus \mathbb{Z}/2\{u\})\{1, t_1\}\{1, t_2, t_2^2\}.$$

This implies the first isomorphism in the theorem. Since there are relations $c_0^2 = 0, 2c_1 = 0, ..., c_1^2 = 0$ in $CH^*(X)$, we get the second isomorphism. □

Next we consider the exceptional group F_4 for $p = 3$. The cohomology $H^*(G/T)$ is given also by Toda-Watanabe [To-Wa]

$$H^*(G/T)_{(3)} \cong \mathbb{Z}_3[t_1, t_2, t_3, t_4, y]/(\rho_1, \rho_2, \rho_3, \rho_4, f_5).$$

Here relations ρ_i is written by using the elementary symmetric function $c_i = \sigma_i(t_1, t_2, t_3, t_4)$, that is,

$$\rho_1 = c_2 - (1/2)c_1^2, \quad \rho_2 = c_4 - c_3c_1 + (1/2)^3c_1^4 - 3y,$$

$$\rho_3 = -c_4c_1^2 + c_3^2, \quad \rho_4 = 3c_4c_1^4 - (1/2)^4c_1^8 + 3y(2^4y + 2^3c_3c_1), \quad f_5 = y^3.$$

The element v_1y is given as ρ_1 by dimensional reason. For $3y, 3y^2$, they are still given in the relations ρ_2, ρ_4. We have proven $\rho_3 = v_1y^2$ in Lemma 9.3.

Let $S(t) = \mathbb{Z}[t_1, ..., t_4]$ and take $\bar{b}_i \in S(t)$ such that

$$H^*(G/T;\mathbb{Z}/p)/(y) \cong \bar{S}(t) \quad with\ \bar{S}(t) = S(t)/(\bar{b}_1, ..., \bar{b}_4).$$

Namely, $b_1 = \rho_1$, $b_2 = \rho_2 + 3y$, $b_3 = \rho_3$, $b_4 = \rho_4 - 3y(2^4y - 2^3c_3c_1)$.

Theorem 10.2. *Let $(G,p)=(F_4,3)$ and $X=\mathbb{G}_k/T_k$. Then there are $b_{i,j}\in S(t)$ for $1\le i,j\le 4$ such that $b_{i,j}=\bar{b}_i\bar{b}_j$ mod(3) and*

$$CH^*(X)\cong S(t)/(3b_1,3b_3,b_{i,j}|1\le i,j\le 4)$$

Remark. For example, we see $b_{2,2}=b_2^2-3b_4$ *mod*(9), since $b_2=3y,\ b_4=3y^2$ *mod*(9).

11. products of groups of type (I)

In this section, we study $R(G_1\times ...\times G_n)$ for groups G_i of type (I). At first we study the case $n=2$. Let $G=G_i$ be of type (I). The (p-localized) torsion index of $G_1\times G_2$ is p^2. Hence we have a field k such that there is a non-trivial $\mathbb{G}_k$ over k for all extensions of index pa (where a is coprime to p) is still nontrivial. Let us write

$$H^*(G_1\times G_2;\mathbb{Z}/p)\cong\mathbb{Z}/p[y_1,y_2]/(y_1^p,y_2^p)\otimes\Lambda(x_1,...,x_\ell).$$

Let $dim(X)=d$ and $f\in H^{2d}(G/T)$ be the fundamental class (e.g. $f=(y_1y_2)^{p-1}s$ for $s\in S(t)$). Consider the degree map $CH^d(X)\overset{\pi_*=deg}{\to}CH^0(pt.)$. When $X=(G_1\times G_2)/T$, the image $Im(\pi^*)=p^2\mathbb{Z}_{(p)}$. Since $\bar{\pi}^*$ is epic, we see $res(CH^d(X))=p^2\mathbb{Z}_{(p)}\{f\}$. So $pf\notin Res_{CH}$.

Lemma 11.1. *Let $G=G_1\times G_2$ for type (I) groups G_i and $X=\mathbb{G}_k/T_k$. Then there are nonzero elements $c_{0,0}$ (resp. $c_{0,1},c_{1,1}$) in $CH^*(X)$ with degree $2d$ (resp. $2d-2(p-1)$, $2d-4(p-1)$) which corresponds to p^2f (resp. pv_1f, v_1^2f) in $\Omega^*(\bar{X})$.*

Proof. First note that $v_1^2y_1^{p-1}y_2^{p-1}=v_1y_1^{p-1}\cdot v_1y_2^{p-1}\in Res_\Omega$. If $v_1f\in Res_\Omega$, then using the Quillen operation $pf\in Res_\Omega$ and this is a contradiction. Thus p^2f,pv_1f,v_1^2f are Ω^*-module generators of Res_Ω. In fact, we have $J\{f\}=Res_\Omega\cap\Omega^*\{f\}$ for the invariant ideal $J=(p^2,pv_1,v_1^2)$ □

Here we assume a strong assumption

$$(*)\quad p^{n-1}y_1^{i_1}...y_n^{i_n}\notin Res_{CH}\quad 1\le i_j\le p-1,\ 1\le j\le n.$$

Proposition 11.2. *Let $G_1,...,G_n$ be groups of type (I) satisfying $(*)$ and let $G=G_1\times ...\times G_n$. Then we have maps*

$$\otimes_{i=1}^n CH^*(\mathbb{G}_{i,k}/B_k)\to CH^*(\mathbb{G}_k/B_k)$$

$$\twoheadrightarrow\otimes_{i=1}^n CH^*(\mathbb{G}_{i,k}/B_k)/(c_0(y_s^i)c_1(y_t^j)=c_1(y_s^i)c_0(y_t^j))$$

with $1\le i,j\le p-1$, $1\le s,t\le n$. Moreover if $\mathbb{G}_k/B_k)$ is generically twisted variety, the first map is also surjective.

Proof. The second surjective follows from considering $\Omega^*(G_k/B_k)$. The first surjection follows from Karpenko's theorem. The relation follows from

$$c_0(y_1^i)c_1(y_2^j) = py_1^i v_1 y_2^j = v_1 y_1^i p y_2^j = c_1(y_1^i)c_0(y_2^j)).$$

□

Next we consider the case that extension of $\mathbb{G}_k$ of degree p splits, namely,

$$(**) \quad py_1^i y_2^j \in Res_{CH} \quad 1 \leq i, j \leq p-1.$$

Then we have $1 \leq deg_v(y_1^i y_2^j) \leq 2$.

At first, we consider the space $X \times X$ for $X = \mathbb{G}_k/T_k$. It is well known that

$$CH^*(X \times X) \twoheadrightarrow CH^*(X \times_k k(X)) = CH^*(X|_{k(X)}) \cong CH^*(\bar{X}).$$

is surjective. In particular $y' = y \otimes 1 + 1 \otimes y$ is in Res_{CH}. We consider the natural map $i : (G \times G)/(T \times T) \to G/T \times G/T$. Let us write $y_1 = i^*(y \otimes 1)$ and $y_2 = i^*(y')$. Then $J(\mathbb{G}_k) = (1, 0)$ for $G = G_1 \times G_1$. Hence we have

Lemma 11.3. *Let $G = G_1 \times G_1$ for a type (I) group G_1 satisfying $(**)$. Let $X = \mathbb{G}_k/T_k$ and $X_1 = (\mathbb{G}_1)_k/T_k$. Then we have*

$$R(\mathbb{G}_k; \mathbb{Z}/p)/p \cong R_2/p,$$

$$CH^*(X) \cong CH^*(R_2) \otimes \mathbb{Z}/p[y'] \otimes S(t)/(\bar{b}_1, ..., \bar{b}_{2\ell})$$

$$\cong CH^*(X_1) \otimes CH^*(\bar{X}_1).$$

This arguments work for general cases. If $G_1 \times G_2$ are both of type (I), then we can take $G_1 \subset G_2$. So we first consider $G = G_2 \times G_2$ and then consider the restriction to $G_1 \times G_2$. Thus we have

Proposition 11.4. *Let $G = G_1 \times ... \times G_s$ for type (I) groups G_i. Suppose that $\mathbb{G}_i$ split over any extension of $a \cdot p$ (a is coprime to p) of k. Then the J-invariant is $J(\mathbb{G}) = (1, 0, ..., 0)$ and*

$$CH^*(\mathbb{G}_k/T)/p \cong CH^*(R_2) \otimes H^*(G/T; \mathbb{Z}/p)/(y_1).$$

Corollary 11.5. *By the same assumption as in the above theorem, we have with $|y_i'| = 2p + 2$*

$$CH^*(\mathbb{G}_k)/p \cong P(y)/(p, y_1) \cong \mathbb{Z}/p[y_2', ..., y_s']/(y_2'^p, ..., y_s'^p).$$

12. Type (II) group

In this section, we consider $CH^*(\mathbb{G})$ for groups of type (II) (e.g., $(G,p)=(E_8,3)$). We have a relation

$$v_1y + v_2y' = 0 \quad mod(I_\infty^2) \ in \ \Omega^*(G_k),$$

which induces from Theorem 6.7

$$(*) \quad v_1y + v_2y' = 0 \quad mod(S(t), I_\infty^2) \ in \ \Omega^*(G_k/T_k).$$

Here it is known $P^py = y'$. So $y \in Res_{CH}$ implies $y' \in Res_{CH}$. In any way we assume that $y \notin Res_{CH}$. Recall that $v_1y \in Res_{\Omega^*}$.

Lemma 12.1. *Let (G,p) be a group of type (II) and $X = \mathbb{G}_k/T_k$. If $y' \in Res_{CH}$, then $v_1y \in Im(res_\Omega) \subset \Omega^*(\bar{X})$.*

Proof. It follows from $(*)$ and $v_2y' \in Im(res_{\Omega^*})$. □

Lemma 12.2. *Let $(G,p) = (E_8,3)$ and $X = \mathbb{G}_k/T_k$. Let $y' \notin Res_{\Omega^*}$. Then $c_1(y)$ exists as the element $v_1y + v_2y'$ in $\Omega^*(\bar{X})$ but v_1y itself (and hence v_2y') is not in $Im(res_\Omega)$.*

Proof. Since $v_1y \in Res_{\Omega^*} \subset gr\Omega^*(\bar{X})$, we have the non zero element $c_0(y)$. In particular, $v_1y + v_2y' \in BP^* \otimes S(t)/I_\infty^2$ from $(*)$. Of course it is non zero in $\Omega^*(\bar{X})$ from (6.3).

On the other hand Garibaldi and Zainouline shows

$$torsion(gr^4_\gamma(X)_{(3)}) \cong \mathbb{Z}/3.$$

(For the filtrations in K-theory, see the last section of this paper.) Since $gr^4_\gamma(X) \to gr^4_{geo(k)}(X)$ is surjective, we know the 3-torsion parts of $gr^4_{geo(k)}(X) \cong CH^2(X)$ is just $\mathbb{Z}/3$ (Proposition 5.4 in [Ga-Za]). If $v_1y \in Im(res_{\Omega^*})$, then the 3-torsion part contains $\mathbb{Z}/3 \oplus \mathbb{Z}/3$. □

Recall the *mod* 3 cohomology of F_4 is given

$$H^*(F_4;\mathbb{Z}/3) \cong \mathbb{Z}/3[y,y']/(y^3,(y')^3) \otimes \Lambda(x_3,x_7,x_{15},x_{19},x_{27},x_{35},x_{39},x_{47}).$$

Here the suffix means its degree, i.e., $|x_i| = i$. By Kono-Mimura [Ko-Mi] the reduced operations act as

$$P^1 : \ x_3 \to x_7, \quad x_{15} \to x_{19}, \quad x_{35} \to x_{39}$$
$$P^3 : \ x_7 \to x_{19}, \quad x_{15} \to x_{27} \to -x_{39}, \quad x_{35} \to x_{47}.$$

We also have the isomorphism

$$grH^*(G/T)_{(3)} \cong \mathbb{Z}_{(3)}[y,y'] \otimes S(t)/(\bar{b}_4, \bar{b}_8, ..., \bar{b}_{48}, f_{24}, f_{60}).$$

More strongly, Nakagawa compute $H^*(G/T)$ (without localization at primes). In his notation

$$H^*(G/T) \cong \mathbb{Z}[\gamma_4, \gamma_{10}, ...] \otimes S(t)/(\rho_4, ...).$$

Here $\gamma_4 = y, \gamma_{10} = y', \rho_2 = \bar{b}_4, ...$ in our notations.

Lemma 12.3. *Let $(G,p) = (E_8, 3)$ and $X = \mathbb{G}_k/T_k$. In $\Omega^*(X)$, there are $b_i \in S(t)$ such that $b_i = \bar{b}_i \ mod(I_\infty)$, and $b_i \neq 0 \in CH^*(X)/3$*

$$v_1 y + v_2 y' = b_4, \quad 3y = b_8, \quad 3y^2 + v_1(y') = b_{16}, \quad 3y' = b_{20},$$
$$3y'y = b_{28}, \quad 3y^2y' + v_1(y')^2 = b_{36}, \quad 3(y')^2 = b_{40}, \quad 3y(y')^2 = b_{48}.$$

Proof. The first relations follows from the preceding lemma. We use cohomology operations $r_{\Delta_1}(v_1) = 3$, $r_{\Delta_2}(v_2) = 3$, and $r_{3\Delta_1}(y) = y'$. We have

$$b_8 = r_{\Delta_1}(b_4) = r_{\Delta_1}(v_1 y + v_2 y') = 3y.$$

Similarly we have b_{20} by acting r_{Δ_2} on b_4.

We need Nakagawa's result such that relations are written

$$\rho_8 = 3\gamma_4^2 + ..., \quad \rho_{18} = -6\gamma_4^2\gamma_{10} + ...$$

Since $r_{\Delta_1}(b_{16}) = b_{20}$, we have $b_{16} = 3y_8^2 + v_1 y'$.

Acting $r_{3\Delta_1}$ and $r_{6\Delta_1}$ on b_{16}, we have b_{28} and b_{40}. By Nakagawa's result above and b_{40}, we have b_{36}. It induces b_{48}. □

Corollary 12.4. *Let $(G,p) = (E_8, 3)$ and $X = \mathbb{G}_k/T_k$ be generically twisted. Then $b_4 b_{48}, b_8 b_{48}$ are non zero in $CH^*(X)/3$.*

Proof. First note that the fundamental class f in $H^*(G/T)$ is represented as $(yy')^2 b$ for some $b \in S(t)$. Since 3 component of the torsion index is 9 and X is generically twisted, we see $3(yy')f \notin res_{CH}$. It follows $3(yy') \notin res_{CH}$ from the theorem by Petrov, Semenov and Zainoulline. Therefore $9(yy')^2, 3v_1(yy')^2$ are BP^*-module generators in Res_Ω. Since $res_\Omega(b_8 b_{48}) = 9(yy')$ and $res_\Omega(b_4 b_{48}) = 3v_1(yy')^2$, we have the corollary. □

13. $p = 2$ case, quadrics

In this section, we rewrite arguments of quadrics by using $deg_v(y)$, $M_n(y)$ and $Q(X)$. Let $h \in CH^1(X)$ be the hyperplane section so that $Res_{CH}(h) = t$. So we use the notation h instead of t. Let X be a quadric of $dim(X) = 2\ell - 1$. From Rost and Toda-Watanabe

$$CH^*(\bar{X}) \cong \mathbb{Z}[h,y]/(h^\ell = 2y, y^2)$$
$$\overset{additively}{\cong} \mathbb{Z}\{1, h, ..., h^{\ell-1}\} \oplus \mathbb{Z}\{y, ..., h^{\ell-1}y\}.$$

Throughout this section, we assume that X is anisotropic. This means $deg_v(h^{\ell-1}y) \geq 1$ since $h^{\ell-1}y$ is the fundamental class of the $(2\ell-1)$-dimensional manifold $\bar{X}$. Thus we get

$$deg_v(h^i) = 0, \quad deg_v(h^i y) \geq 1 \quad for\ all\ 0 \leq i \leq \ell - 1.$$

Recall the quotient $Q(X)$ of $CH^*(X)$

$$Q(X) = \oplus_{0\leq i\leq \ell-1}(M_0(h^i) \oplus M_{d_i}(h^i y)) \quad with\ d_i = deg_v(h^i y).$$

Since $h \in Res_{CH}$, it is immediate $d_0 \leq d_1 \leq ... \leq d_{\ell-1}$.

We recall that

$$M_0(h^i) \oplus M_{d_i}(h^i y) \cong \mathbb{Z}\{h^i, c_0(h^i y)\} \oplus \mathbb{Z}/2\{c_1(h^i y), ..., c_{d_i-1}(h^i y)\}.$$

We note $c_0(h^i y) = 2h^i y = h^{\ell+i}$. Hence we can write

$$Q(X) \cong \mathbb{Z}[h]/(h^{2\ell}) \oplus \oplus_{0\leq i\leq \ell-1}\mathbb{Z}/2\{c_1(h^i y), ..., c_{d_i-1}(h^i y)\}.$$

Note that the Rost motive is isomorphic to

$$R_{d_i}[2J_i](J_i) \cong \mathbb{Z}\{h^{J_i}, c_0(h^i y)\} \oplus \mathbb{Z}/2\{c_1(h^i y), ..., c_{d_i-1}(h^i y)\}$$

where

$$J_i = 1/2(|h^i y| - |R_{d_i}|) = i + 2^\ell - 1 - (2^{d_i} - 1) = i + (2^\ell - 2^{d_i}).$$

Therefore if the motive $M(X)$ is a sum of Rost motives (e.g., excellent quadrics), then the above quotient ring is $CH^*(X)$ itself. In particular the set $\{J_0, ..., J_{\ell-1}\} = \{0, 1, ..., \ell - 1\}$.

Theorem 13.1. *The Chow ring $CH^*(X)$ has the quotient ring*

$$Q(X) \cong \mathbb{Z}[h]/(h^{2\ell}) \oplus_{j=0}^{s} \mathbb{Z}/2[h]/(h^{\ell-f_j})\{u_j\}$$

where f_j is the smallest f such that $v_j h^f y \in Res_\Omega$ (i.e., $deg_v(h^f y) = j + 1$), and $u_j = c_j(h^{f_j} y) = v_j h^{f_j} y$, and so $s = deg_v(h^{\ell-1} y) - 1$.

Proof. This is immediate by changing order of sums such that

$$Q(X) \cong \oplus I_{d_i}\{h^i y\} = \oplus_{i,j}(v_j h^i y) = \oplus h^{i-f_j}(v_j h^{f_j}) = \oplus h^{i-f_j} u_j.$$

□

Therefore the Chow ring of each excellent quadric is isomorphic to the above quotient ring $Q(X)$. For example, we consider when $\ell = 2^n - 1$ and $k = \mathbb{R}$ (the field of real numbers). Recall that $\rho = (-1) \in K_1^M(\mathbb{R})/2$ and $\phi_{\rho^{n+1}} = \langle\langle -1, ..., -1\rangle\rangle$ is the $(n+1)$-th Pfister form associated to ρ^{n+1}. Let q' is the subform of codimension 1, that is, q' is the maximal neighbor of the $(n+1)$-th Pfister form $\phi_{\rho^{n+1}}$.

Lemma 13.2. *([Ya3]) Let $k = \mathbb{R}$ and $X = X_{q'} = \mathbb{G}_k/P_k$ be the maximal neighbor of the $(n+1)$ Pfister quadric. Then $deg_v(yh^i) = n$ for all $0 \leq i \leq \ell - 1$. Hence we have the ring isomorphism*

$$CH^*(X) \cong \mathbb{Z}[h]/(h^{2^{n+1}-2}) \oplus \mathbb{Z}/2[h]/(h^{2^n-1})\{u_1, ..., u_{n-1}\}$$

where $u_i = c_i(y) = v_i y \in \Omega^*(\bar{X})$ so that $u_i u_j = 0$.

Lemma 13.3. *([Ya3]) Let $k = \mathbb{R}$, $\ell = 2^{n-1}$, and $X = \mathbb{G}_k/P'_k$ be the minimal neighbor of the $(n+1)$ Pfister quadric. Then $deg_v(vh^{\ell-1}) = n$ and $deg_v(yh^i) = n-1$ for all $0 \leq i \leq \ell - 2$. Hence we have the ring isomorphisms*

$$CH^*(X) \cong \mathbb{Z}[h]/(h^{2^n}) \oplus \mathbb{Z}/2[h]/(h^{2^{n-1}})\{u'_1, ..., u'_{n-2}\} \oplus \mathbb{Z}/2\{u'_{n-1}\}$$

where $u'_i = c_i(y)$ for $1 \leq i \leq n-2$ and $u'_{n-1} = c_{n-1}(h^{2^{n-1}-1}y)$.

Let $f : X \to Y$ be a map of anisotropic quadrics. Suppose $f^*(h_Y) \neq 0 \in CH^2(\bar{X})/2$ (i.e. $f^*(h_Y) = h_X$). Let y_Y be the ring generator of $CH^*(\bar{Y})$ such that $2y_Y = h^{\ell_Y}$ for $\ell = 1/2(dim(Y)-1)$. Similarly define the ring generator y_X in $CH^*(\bar{X})$. Then $f^*(y_Y) = h^{\ell_Y - \ell_X} y_X$, and so $dim(X) \leq dim(Y)$. Thus we see

(1) $f^* : CH^*(\bar{Y}) \to CH^*(\bar{X})$ is injective for $* \leq dim(X)$.
(2) each $h^j y_X \notin Res_{CH}$ for $0 \leq j \leq \ell_X - 1$.

From Corollary 3.7, we have

Theorem 13.4. *Let $f : X \to Y$ be a map of anisotropic quadrics such that $f : CH^2(\bar{Y})/2 \to CH^2(\bar{X})/2$ is non zero. Then the induced map $Q^{\leq dim(X)}(Y) \to Q(X)$ is injective.*

Corollary 13.5. *If X is a neighbor of the $n+1$-Pfister quadric, then $deg_v(X) = n$.*

Example. Let X_{min} (resp. X_{max}) be the minimal (resp. maximal) neighbor of the $(n+1)$-Pfister quadric and $e : X_{min} \subset X_{max}$ is the embedding. Note $dim(X_{min}) = 2^n - 1$ and we see

$$Q^{\leq 2^n-1}(X_{max}) \cong \mathbb{Z}[h]/(h^{2^n}) \oplus \mathbb{Z}/2\{u_1, ..., u_{n-1}\}$$

from Lemma 14.3. Hence we see that it injects into $Q(X_{min})$, by $e^*(u_i) = h^{2^{n-1}-1}u'_i$ for $i \leq n-2$ and $e^*(u_{n-1}) = u'_{n-1}$.

Corollary 13.6. *Let Y be an anisotropic quadric and $X = Y \cap H^{2d}$ for $2d$-dimensional plane H^{2d}, and $e : X \subset Y$ be the induced embedding. Then the Gysin map $e_* : Q(X) \to Q(Y)$ is injective. Let us write by $f_j(X)$ (resp. $f_j(Y)$) the smallest number f such that $v_j h^f y_X \in Res_\Omega$ (resp. $v_j h^f y_Y \in Res_\Omega$). Then we have*

$$f_j(Y) - d \leq f_j(X) \leq f_j(Y) + d.$$

Proof. First note that $e^*(y_Y) = h^d y_X$. Note $e_*(1) = h^{2d}$ and $e_*(y_X) = h^d y_Y$. Let $f = f_j(Y)$. Then $v_j h^f y_Y \in Res_\Omega$ and hence

$$e^*(v_j h^f y_Y) = v_j h^{f+d} y_X \in Res_\Omega.$$

This means $f_j(X) \leq f_j(Y) + d$. If $v_j h^g y_X \in Res_\Omega$ then $e_*(v_j h^g y_X) = v_j h^{g+d} y_Y \in Res_\Omega$. This fact implies $f_j(Y) \leq f_j(X) + d$. □

14. The orthogonal group $SO(m)$ and $p = 2$

We consider the orthogonal groups $G = SO(m)$ and $p = 2$. Of course $SO(m)$ is not simply connected, but it is well known that $SO(m)/T \cong Spin(m)/T'$ for the spin group $Spin(m)$ and its maximal torus T'. Hence we consider $SO(m)/T$ instead of $Spin(m)/T'$. The mod 2-cohomology is written as (see for example [Ni])

$$grH^*(SO(m); \mathbb{Z}/2) \cong \Lambda(x_1, x_2, ..., x_{m-1})$$

where $|x_i| = i$, and the multiplications are given by $x_s^2 = x_{2s}$. We write $y_{2(odd)} = x_{odd}^2$. Hence we can write

$$H^*(SO(m); \mathbb{Z}/2) \cong P(y) \otimes \Lambda(x_1, x_3, ...x_{\bar{m}})$$

$$with \quad P(y) = \otimes_{i=0}^{t} \mathbb{Z}/2[y_{4i+2}]/(y_{4i+2}^{2^{s(i)}}), \quad grP(y) \cong \Lambda(x_2, x_4, ...x_{m'})$$

where $2 \leq 4i + 2 \leq m - 1$, and $s(i)$ is the smallest number such that $m \leq 2^{s(i)}(4i + 2)$, t is the largest number such that $4t + 1 \leq m - 1$, and where $\bar{m} = m - 1$ (resp.$\bar{m} = m - 2$) if m is even (resp. odd) and where $m' = m - 2$ (resp.$\bar{m} = m - 1$) if m is even (resp. odd). The index means its degree, namely $|y_j| = j, |x_k| = k$.

The Steenrod operation is given as

$$Sq^k(x_i) = \binom{i}{k}(x_{i+k}).$$

The Q_i-operations are given by Nishimoto [Ni]

$$Q_n x_{odd} = x_{odd+|Q_n|}, \qquad Q_n x_{even} = Q_n y_{even} = 0$$

Relations in $\Omega^*(SO(m))$ are given by

$$\sum_n v_n Q_n(x_{odd}) = \sum_n v_n x_{odd+|Q_n|} = 0 \qquad mod(I_\infty^2).$$

For example, the relation in $\Omega^*(SO(m))/I_\infty^2$ starting with $2y_6$ is given by

$$2y_6 + v_1 y_2^4 + v_2 y_6^2 + v_3 y_{10}^2 + ... = 0.$$

Theorem 14.1. *([Ya1]) There is an $\Omega^*_{(2)}$-algebra isomorphism*

$$\Omega^*(SO(m))/I_\infty^2 \cong \Omega^* \otimes_i \mathbb{Z}[y_{4i+2}]/(R, I_\infty^2)$$

where $2 \leq 4i + 2 \leq m - 1$ and R is the relations which start with

$$y_{4i+2}^{2^{s(i)}}, \quad 2y_{4i+2}, \quad v_1 y_{4i'+2} \ for \ i' \neq 0.$$

For ease of arguments, we only consider the case $G = SO(odd)$. Let $G = SO(2\ell + 1)$ and $P = SO(2\ell - 1) \times SO(2)$. Then it is well known [To-Wa]

Lemma 14.2. $H^*(G/P) \cong \mathbb{Z}[t,y]/(t^\ell - 2y, y^2) \quad |y| = 2\ell.$

By Toda-Watanabe [To-Wa], we also know

Theorem 14.3. *(see the proof of Theorem 2.1 in [To-Wa])*

$$H^*(G/T) \cong \mathbb{Z}[t_i, y_{2i}, t_\ell, y]/(c_i - 2y_{2i}, J_{2i}, t_\ell^\ell - 2y, y^2)$$

where $1 \leq i \leq \ell - 1$, $c_i = \sigma(t_1, ..., t_\ell)$ *and*

$$J_{2i} = 1/4(\sum_{j=0}^{2i}(-1)^j c_j c_{2i-j}) = y_{4i} - \sum_{0<j<2i}(-1)^j y_{2j} y_{4i-2j}.$$

Corollary 14.4. *We have* $grH^*(G/T) \cong grH^*(G/P) \otimes A$, *where*

$$A = \mathbb{Z}[t_i, y_i]/(c_i' - 2y_i, J_{2i} | 1 \leq i \leq \ell - 1)$$

with $c_i' = \sigma(t_1, ..., t_{\ell-1})$.

Let us write $P(y)' = \otimes_{i<2^{n-1}-1}\mathbb{Z}[y_{4i+2}]/(y^{2^{s_i}})$ such that $P(y) = P(y)' \otimes \mathbb{Z}[y]/(y^2)$.

We consider the following assumption

$$(*) \quad y_i \in Res_{CH} \quad for\ all\ 1 \leq i \leq \ell - 1.$$

Lemma 14.5. *Suppose* $(*)$ *and* $\ell \neq 2^n - 1$ *for* $n > 0$. *Then* $y_\ell \in Res_{CH}$.

Proof. We see that only $y_{2(2^n-1)}$ is not the target of Steenrod operation. Recall

$$Sq^{2k}(y_{2i}) = \binom{i}{k} y_{2(i+k)}.$$

It is well known that if $i = \sum i_s 2^s$ and $k = \sum k_s 2^s$, then (in $mod(2)$)

$$\binom{i}{k} = \binom{i_m}{k_m} ... \binom{i_s}{k_s} ... \binom{i_0}{k_0}.$$

Note that if $i = 2^n - 1$, then all $i_s = 1$ (for $s < n$). Otherwise there is s such that $i_s = 1$ but $i_{s-1} = 0$. Take $k = 2^{s-1}$ and $i' = i - 2^{s-1}$. Then $i' + k = i$ and

$$\binom{i'}{k} = \binom{i_m = 1}{0} ... \binom{i_s' = 0}{k_s = 0} \binom{1}{1} ... \binom{i_0}{0} \neq 0.$$

This means $Sq^{2k}(y_{2i'}) = y_{2i}$ if $i \neq 2^n - 1$. □

Lemma 14.6. *Suppose* $(*)$ *and* $x \notin Res_{CH}$. *Then if* $\ell = 2^{n+1} - 1$, *then* $deg_v(y) \geq n + 1$.

Proof. By using the result about $Q_i x_j$ by Nishimoto, we have the relation with $mod(I_\infty^2)$ in $\Omega^*(G)$

$$2y_{2\ell-2(2^n-1)} + ... + v_{n-1}y_{2\ell-2(2^n-2^{n-1})} + v_n y.$$

Hence in $\Omega^*(G/T)$, we know

$$v_n y = \sum v(I) y_I t_I \quad with\ v(I) \in I_\infty/(I_\infty^2), t_I \in S(t),$$

and $y_I \in P(y)' \subset A$, which are in Res_{CH} by the assumption $(*)$. Therefore $v_n y \in Res_\Omega$. □

Now we recall an argument of quadrics. Let $m = 2\ell + 1$. and let us write the quadratic form $q(x)$ defined by

$$q(x_1, ..., x_m) = x_1 x_2 + ... + x_{m-2}x_{m-1} + x_m^2$$

and the projective quadric X_q defined by the quadratic form q. Then it is well known that (in fact $SO(m)$ acts on the affine quadric in $\mathbb{A}^m - 0$)

$$X_q \cong SO(m)/(SO(m-2) \times SO(2)).$$

Let $G = SO(m)$ and $P = SO(m-2) \times SO(2)$. Then the quadric q is always split over k and we know $CH^*(G_k/P_k) \cong CH^*(X_q)$.

In particular we consider the case $m = 2^{n+1} - 1$ (hence $\ell = 2^n - 1$). Recall $\rho = \{-1\} \in K_1^M(k) = k^*$. We consider the field k such that

$$(**) \qquad 0 \neq \rho^{n+1} \in K_{n+1}^M(k)/2.$$

Throughout this section, we assume the condition $(**)$. Define the quadratic form q' by $q'(x_1, ..., x_m) = x_1^2 + ... + x_m^2$. Then this q' is the maximal neighbor of the $(n+1)$-th Pfister form associated to ρ^{n+1}. Of course $q|_{\bar{k}} = q'|_{\bar{k}}$ and we can identify $\mathbb{G}_k/P_k \cong X_{q'}$. A quadric is generically split if it is a Pfister quadric or its maximal neighbor. Moreover we know that $(*)$ is satisfied this case (i.e. the J-invariant is $(0, ..., 0, 1)$ see 7.5 in [Pe-Se-Za]). From Lemma 14.5 and Theorem 8.4 (the theorem by Petrov, Semenov and Zainoulline [Pe-Se-Za]), we know

Proposition 14.7. *Let $m = 2^{n+1} - 1$ and $\mathbb{G}_k/P_k \cong X_{q'}$ for the maximal neighbor q' of the $n+1$-th Pfister form. Then we have*

$$CH^*(\mathbb{G}_k/T_k) \cong CH^*(X_{q'}) \otimes A.$$

As a corollary, we see that t_i, y_i are all in $CH^*(\mathbb{G}/T_k)$. Hence $CH^*(\mathbb{G}/T_k)$ is multiplicatively generated by t_i, y_i, t and $u_1, ..., u_{n-1}$.

Corollary 14.8. *(Theorem 7.6 in [Ya5]) Assume $(**)$ and $m = 2^{n+1} - 1$. Then $CH^*(\mathbb{G}_k)_{(2)} \cong P(y)'/(2)$.*

At last of this section, we consider the case $\bar{X} = X(\mathbb{C}) = G/P$ with

$$G = SO(2\ell+1) \quad and \quad P = U(\ell).$$

From the cofibering $SO(2\ell+1) \to \bar{X} \to BU(\ell)$, we have the spectral sequence

$$E_2^{*,*} \cong H^*(SO(2\ell+1); \mathbb{Z}/2) \otimes H^*(BU(\ell)) \Longrightarrow H^*(\bar{X}; \mathbb{Z}/2).$$

Here $H^*(BU(\ell) \cong \mathbb{Z}[c_1, ..., c_\ell]$ and the differential is given as $d_{2i}(x_{2i-1}) = c_i$. Hence

$$H^*(\bar{X}; \mathbb{Z}/2) \cong H^*(SO(2\ell+1); \mathbb{Z}/2)/(x_1, x_3, ..., x_{2\ell-1}) \cong P(y).$$

Let Q be a quadric of dimension $2\ell-1$, and $G(\ell, Q)$ the grassmanian of projective spaces of dimension $\ell - 1$ on Q. Then $G(\ell, Q)|_{\bar{k}} = \bar{X} = SO(2\ell+1)/U(\ell)$ and so

$$CH^*(G(\ell, Q)|_{\bar{k}}) \cong H^*(SO(2\ell+1)/U(\ell)).$$

This variety $X = G(\ell, Q)$ is studied by Vishik in [Vi1]. He shows the case G/B also. In his notation, $F(\ell, Q)$ is the complete flag variety , i.e., $F(\ell, Q) = \mathbb{G}_k/B_k$ for $G = SO(2\ell+1)$.

Theorem 14.9. *(Vishik Theorem 5.1 in[Vi1]) We have the isomorphism for* $X = \mathbb{G}_k/B_k = F(\ell, Q)$

$$CH^*(X)/2 \cong CH^*(G(\ell, Q)/2 \otimes H^*(U(\ell)/T)$$

where $H^*(U(\ell)/T) \cong \mathbb{Z}[t_1, ..., t_\ell]/(c_1, ..., c_\ell)$.

Note that the integral cohomology (hence the Chow ring) $H^*(\bar{X})$ still given by Toda-Watanabe (Theorem 14.3). The main theorem in [Vi] is the following. Recall $grP(y) \cong \Lambda(x_2, x_4, ..., x_{2\ell})$.

Theorem 14.10. *(Vishik Theorem 5.8 in [Vi1]) Let* $(i_1, ..., i_s) \subset (1, 2, ..., \ell)$ *be the maximal subset with each* $x_{2i_s} \in Res_{CH}$. *Let* $y \in CH^*(G(\ell), Q)|_{\bar{k}}) \cong P(y)$. *Then*

$$y \in Res_{CH} \quad if\ and\ only\ if \quad y \in \Lambda(x_{2i_1}, ..., x_{2i_s}).$$

Remark. Vishik defined the J-invariant as $J(G(\ell, Q)) = \{i_s | x_{2i_s} \in Res_{CH}\}$ in [Vi1].

If $deg_v(y) \leq n$, then $deg_v(y^2) \leq n$, since $Sq^{|y|}(y) = y^2$. It seems that $deg_v(y), deg_v(y') \leq n$ does not implies $deg_v(yy') \leq n$. However we consider its converse in the following sense

Proposition 14.11. *Let* $(i_1, ..., i_s) \subset (1, ..., \ell)$. *If* $deg_v(x_{2i_1}...x_{2i_s}) \leq n$, *then* $deg_v(x_{2i_j}) \leq n$ *for each* $1 \leq j \leq s$.

The proof of this proposition is given by the $deg_v(-)$ version of the proof of the above Vishik's theorem. Vishik use the Schubert cells. In his notation $\bar{z}_i = y_{4(i-1)+2}$. It is well known

$$\bar{z}_I \cdot \bar{z}_j = \bar{z}_{I\cup j} + \sum_{i\in I} \bar{z}_{(I-i)\cup(i+j)} \quad for\ I \subset (1....,\ell).$$

Let $\pi : F(\ell, Q) \to F(\ell-1, Q)$ be the natural projection between full flag varieties. Vishik shows (Proposition 5.4 in [Vi1])

$$\pi^*\pi_*(\bar{z}_I) = \sum_{i\in I} \bar{z}_{I-i} \cdot \pi^*\pi_*(\bar{z}_i).$$

The above theorem is proved as follow (proof of Lemma 5.9 in [Vi1]). Let $I = (i_1, ..., i_s) \subset (1, ..., \ell)$. Denote the operation $\pi^*\pi_*$ as D. Then $D(\bar{z}) = \sum_{1\le i\le \ell} d_i(\bar{z}) \cdot D(\bar{z}_i)$ where $d_i(\bar{z}) \in CH^*(G(\ell, Q)|_{\bar{k}})/2$ and $D(z_i) \in CH^*(\bar{U}(\ell)/T)$. Then we can see

$$d_{i_s}...d_{i_2}(\bar{z}_I) = z_{i_1}.$$

Thus $z_I \in Res_{CH}$, then so is z_{i_1}.

For the proof of the proposition, let $v_n z_I \in Res_\Omega$. Then we can see

$$d_{i_s}...d_{i_2}(v_n\bar{z}_I) = v_n z_{i_1} \in Res\ mod(I_\infty^2).$$

15. FILTRATIONS IN K-THEORY

We consider the cases that

$$(1.1) \quad K^0_{alg(k)}(X) \cong K^0_{alg(\bar{k})}(\bar{X}) \cong K^0_{top}(X(\mathbb{C}))$$

where $K^0_{alg(k)}(X)$ (resp. $K^0_{top}(X(\mathbb{C}))$) is the algebraic (resp. topological) K-theory generated by algebraic k-bundles (resp. complex bundles) over X (resp. $X(\mathbb{C})$). In this assumption, we study the typical filtrations

$$F^i_\gamma(X) \ \subset\ F^i_{geo(k)}(X) \subset F^i_{geo(\bar{k})}(\bar{X}) \subset\ F^i_{top}(X(\mathbb{C}))$$

namely, the gamma and the geometric filtrations defined by Grothendieck [Gr], and the topological filtration defined by Atiyah [At]. Namely, we study induced maps of associated rings

$$gr^*_\gamma(X) \to gr^*_{geo(k)}(X) \to gr_{geo(\bar{k})}(\bar{X}) \to gr^*_{top}(X(\mathbb{C})).$$

Note that $gr_\gamma(X) \cong gr_\gamma(X(\mathbb{C}))$ by the above isomorphisms of K-theories and definition of the γ-filtration ([To3,Ya7]).

Atiyah showed that $gr^*_{top}(X(\mathbb{C}))$ is isomorphic to the infitite term $E^{*,0}_\infty$ of the AHss converging to K-theory $K^*(X(\mathbb{C}))$. Moreover he showed that $gr^*_{top}(X(\mathbb{C})) \cong gr^*_\gamma(X)$ if and only if $E^{*,0}_\infty$ is generated by Chern classes in $H^*(X(\mathbb{C}))$. We will see that similar facts hold for

$gr^*_{geo}(X)$. Namely, $gr^{2*}_{geo}(X) \cong AE^{2*,*,0}_\infty$ of the motivic AHss converging to motivic K-theory $AK^{*,*'}(X)$ ([Ya7]). Moreover we show that $gr^*_{geo}(X) \cong gr^*_\gamma(X)$ if and only if $AE^{2*,*,0}_\infty$ is generated by Chern classes in the Chow ring $CH^*(X) \cong H^{2*,*}(X)$.

We consider the following assumption for X.

Assumption $(*)$. There is an isomorphism of motives

$$M(X) \cong R_n \otimes A(X) \quad with\ A(X) \cong \oplus_s \mathbb{T}^{i_s}$$

where $\mathbb{T}$ is the k-Tate module. From (the case $n = 1$ of) Lemma 7.3, we have

$$K^0_{alg}(X) \cong K^0_{alg}(X|_{\bar{k}}) \cong K^0_{top}(X(\mathbb{C})).$$

Note that $H^*(G/T)$ is torsion free, and we have

$$gr^*_{geo(\bar{k})}(G_k/T_k) \cong gr^*_{top}(G/T) \cong H^*(G/T).$$

Theorem 15.1. *Let $M(X) = R_2 \otimes A$ for a sum A of Tate modules. If $CH^*(X)$ is generated by Chern classes, then we have the isomorphisms*

$$gr^{2*}_\gamma(X(\mathbb{C}) \cong gr^{2*}_\gamma(X) \cong gr^{2*}_{geo(k)}(X) \cong CH^*(X).$$

Proof. The second isomorphism follows from the fact that $CH^*(X)$ is generated by Chern classes.

We note that

$$gr\Omega^*(X) \cong (\Omega^*\{1\} \oplus (\Omega^*/I_1\{c_1\} \oplus \Omega^*/I_0\{c_0\})[y]/(y^{p-1})) \otimes A.$$

Here $\Omega^*/I_1 = \Omega^*/(p)$ and $\Omega^*/I_0 = \Omega^*$, which are of course v_1-torsion free. Recall $A\tilde{K}(1)^*(X) \cong \Omega^*(X) \otimes_{BP^*} \mathbb{Z}[v_1, v_1^{-1}]$. Hence

$$grA\tilde{K}(1)^*(R_2) \cong \Omega^*(R_a) \otimes_{BP^*} \mathbb{Z}[v_1, v_1^{-1}]$$

$$\cong (\mathbb{Z}_{(p)} \oplus (\mathbb{Z}_{(p)}\{c_0\} \oplus \mathbb{Z}/p\{c_1\})[y]/(y^{p-1})) \otimes \mathbb{Z}[v_1, v_1^{-1}]$$

$$\cong CH^*(R_2) \otimes \mathbb{Z}[v_1, v_1^{-1}]. \quad (*)$$

On the other hand, from AHss converging $A\tilde{K}(1)^*(X)$, it is known

$$A\tilde{K}(1)^*(X) \cong \tilde{K}(1)^* \otimes CH^*(X)/I$$

for some ideal $I \subset CH^*(X)$ (note the differential $d_* = 0$ on $CH^*(X) = H^{2*,*}(X)$). However from $(*)$ above, we see $I = 0$. This implies $gr_{geo(k)}(X) \cong CH^*(X)$. □

From Theorm 9.7 (in fact t_i is represented by a first Chen class), we have

Corollary 15.2. *Let G is a group of type (I) and let $X = \mathbb{G}_k/T_k$. Then we have the isomorphisms in the above theorem.*

Remark. When $M(X) = R_n \otimes A$ for $n \geq 3$, we see

$$gr^{2*}_{geo(k)}(X) \cong CH^*(X)/I \quad with\ I = \mathbb{Z}/p\{c_2, ..., c_{n-1}\}[y]/(y^{p-1}) \otimes A.$$

When $\mathbb{G}_k$ is generically twisted G_k-torsor, by Karpenko, the natural map $CH^*(BB_k) \to CH^*(X)$ for $X = \mathbb{G}_k/B_k)$ is surjective. Hence $CH^*(X)$ generated by the first Chern classes t_i. Therefore we have

Corollary 15.3. *Let $\mathbb{G}_k$ be generically twisted G_k-torsor and $X = \mathbb{G}_k/B_k$. Then*

$$gr_\gamma(X) \cong gr_{geo}(X) \cong CH^*(X)/I$$

for some ideal $I \subset CH^*(X)$. Moreover if $K^0_{alg(k)}(X) \cong K^0_{top}(X(\mathbb{C}))$, then $gr_\gamma(X(\mathbb{C})) \cong CH^*(X)/I$.

References

[Ga-Za] S. Garibaldi and K. Zainoulline. The γ-filtration and the Rost invariant. *J. Reine Angew. Math.* **696** (2014), 225-244.

[Gr] A. Grothendieck. Torsion homologique et sections rationnelles. *Sem. C.Chevalley, ENS 1958,expose 5, Secreatariat Math., IHP, Paris* (1958).

[Ha] M.Hazewinkel, Formal groups and applications, *Pure and Applied Math. 78, Academic Press Inc.* (1978), xxii+573pp.

[Ho] L. Hodgkin. The equivariant Künneth theorem in K-theory, Topics in K-theory, Two independent contributions. Lect. Note. Springer, Berlin, **496** (1975), 1-101.

[Ka] V. G. Kac. Torsion in cohomology of compact Lie groups and Chow rings of reductive algebraic groups. *Invent. Math.* **80** (1985), 69-79.

[Kar] N. Karpenko. Chow groups of some generically twisted flag variety. *www.ualberta.ca/"karpenko.* (2016).

[Ko-Mi] A. Kono and M. Mimura. Cohomology operations and the Hopf algebra structures of compact exceptional Lie groups E_7 and E_8. *Proc London Math. Soc.* **35** (1977), 345-358.

[L-M 1] M. Levine and F. Morel. Cobordisme algébrique I. *C. R. Acad. Sci. Paris* **332** (2001), 723-728.

[L-M 2] M. Levine and F. Morel. Cobordisme algébrique II. *C. R. Acad. Sci. Paris* **332** (2001), 815-820.

[Me-Su] A. Merkurjev and A. Suslin. *Motivic cohomology of the simplicial motive of a Rost variety.* J. Pure and Appl. Algebra **214** (2010), 2017-2026.

[Mi-Ni] M. Mimura and T. Nishimoto. Hopf algebra structure of Morava K-theory of exceptional Lie groups. *Contem. Math.* **293** (2002), 195-231.

[Mi-To] M. Mimura and H. Toda, *Topology of Lie groups*, I and II, Translations of Math. Monographs, Amer. Math. Soc, **91** (1991).

[Na] M. Nakagawa. The integral cohomology ring of E_8/T. *Proc. Japan* (20), 383-394.

[Ni] T. Nishimoto. Higher torsion in Morava K-thoeory of $SO(m)$ and $Spin(m)$. *J.Math.Soc. Japan.* **52** (2001), 383-394.

[Or-Vi-Vo] D.Orlov,A.Vishik and V.Voevodsky. An exact sequence for Milnor's K-theory with applications to quadric forms. *Ann. of Math.* **165** (2007) 1-13.

[Pe-Se-Za] V.Petrov, N.Semenov and K.Zainoulline. J-Invariant of linear algebraic groups. *Ann. Scient. Ec. Norm. Sup.* **41**, (2008) 1023-1053.

[Ra] D.Ravenel. Complex cobordism and stable homotopy groups of spheres. *Pure and Applied Mathematics, 121. Academic Press* (1986).

[Ro1] M.Rost. Some new results on Chowgroups of quadrics. *preprint* (1990).

[Ro2] M.Rost. On the basic correspondence of a splitting variety. *preprint* (2006)

[Sm-Vi] A. Smirnov and A. Vishik.. Subtle characteristic classes. *arXiv : 1401.6661v1 [math.AG]* (2014)

[Su-Jo] A.Suslin and S.Joukhovistski. Norm Variety. *J.Pure and Appl. Algebra 206* (2006) 245-276.

[To1] B. Totaro. The torsion index of E_8 and the other groups. *Duke Math. J.* **299** (2005), 219-248.

[To2] B. Totaro. The torsion index of the spin groups. *Duke Math. J.* **299** (2005), 249-290.

[To3] B. Totaro, Group cohomology and algebraic cycles, *Cambridge tracts in Math. (Cambridge Univ. Press)* **204** (2014).

[Tod] H.Toda. On the cohomolgy ring of some homogeneous spaces. *J. Math. Kyoto Univ.* **15** (1975), 185-199.

[To-Wa] H.Toda and T.Watanabe. The integral cohomology ring of F_4/T and E_4/T. *J. Math. Kyoto Univ.* **14** (1974), 257-286.

[Vi] A. Vishik. Algebraic cobordism as a module over the Lazard ring. arXiv:1409;0156v1. (math.AG) (2012)

[Vi-Ya] A.Vishik and N.Yagita. Algebraic cobordisms of a Pfister quadric. *London Math. Soc.* **76**(2007), 586-604.

[Vi-Za] A.Vishik and K.Zainoulline. Motivic splitting lemma. *Doc. Math.* **13** (2008), 81-96.

[Vo1] V. Voevodsky, The Milnor conjecture, *www.math.uiuc.edu/K-theory/0170* (1996).

[Vo2] V. Voevodsky. Motivic cohomology with $\mathbb{Z}/2$ coefficient. *Publ. Math. IHES* **98** (2003), 59-104.

[Vo3] V.Voevodsky. Reduced power operations in motivic cohomology. *Publ.Math. IHES* **98** (2003),1-57.

[Vo4] V.Voevodsky. On motivic cohomology with $\mathbb{Z}/l$-coefficients. *Ann. of Math.* **174** (2011), 401-438.

[Ya1] N. Yagita. Algebraic cobordism of simply connected Lie groups. *Math.Proc.Camb.Phil.Soc.* **139** (2005), 243-260.

[Ya2] N. Yagita. Applications of Atiyah-Hirzebruch spectral sequence for motivic cobordism. *Proc. London Math. Soc.* **90** (2005) 783-816.

[Ya3] N. Yagita. Chow rings of excellent quadrics. *J. Pure and Appl. Algebra* **212** (2008), 2440-2449.

[Ya4] N. Yagita. Motivic cohomology of quadrics and coniveau spectral sequence. *J. K-theory* **6** (2010), 547-589.

[Ya5] N. Yagita. Note on Chow rings of nontrivial G-torsors over a field. *Kodai Math. J.* **34** (2011), 446-463.

[Ya6] N. Yagita. Algebraic BP-theory and norm varieties. *Hokkaido Math. J.* **41** (2012), 275-316.

[Ya7] N. Yagita. Note on the filtrations of the K-theory. *Kodai Math. J.* **38** (2015), 172-200.

FACULTY OF EDUCATION, IBARAKI UNIVERSITY, MITO, IBARAKI, JAPAN
E-mail address: nobuaki.yagita.math@vc.ibaraki.ac.jp,